arXiv:1302.6727v1 [math.PR] 27 Feb 2013

Frozen percolation in two dimensions

Demeter Kiss*

28th February 2013

Abstract

Aldous [2] introduced a modification of the bond percolation process on the binary tree where clusters stop growing (freeze) as soon as they become infinite. We investigate the site version of this process on the triangular lattice where clusters freeze as soon as they reach L^∞ diameter at least N for some parameter N. We show that, informally speaking, in the limit $N \to \infty$, the clusters only freeze in the critical window of site percolation on the triangular lattice. Hence the fraction of vertices that eventually (i. e. at time 1) are in a frozen cluster tends to 0 as N goes to infinity. We also show that the diameter of the open cluster at time 1 of a given vertex is, with high probability, smaller than N but of order N. This shows that the process on the triangular lattice has a behaviour quite different from Aldous' process. We also indicate which modifications have to be made to adapt the proofs to the case of the N-parameter frozen bond percolation process on the square lattice. This extends our results to the square lattice and answers the questions posed by van den Berg, de Lima and Nolin in [16].

Keywords and phrases: frozen cluster, critical percolation, near critical percolation, correlation length.
AMS 2010 classifications. Primary: 60K35; Secondary: 82B43.

Contents

1 Introduction **2**

2 Preliminary results on near critical percolation **5**
2.1 Correlation inequalities 5
2.2 Mixed arm events, critical arm exponents 6
2.3 Near-critical scaling and correlation length 8

3 Proof of the main results **10**
3.1 Proof of Theorem 1.3 12
3.2 Proof of Corollary 1.6 13

4 Proof of Proposition 3.6 **14**
4.1 Notation 14
4.2 Thick paths 14

5 Proof of Proposition 3.5 **16**
5.1 Lowest point of the lowest crossing in parallelograms 16
5.2 Lowest point of the lowest crossing in regular regions 19
5.3 The diameter of the active clusters close to time 1/2 21

A Appendix **31**
A.1 Winding number of arms 31
A.2 Existence of long thick paths in nice regions 33

---

*Centrum Wiskunde & Informatica, Amsterdam, research supported by NWO; D.Kiss@cwi.nl

1 Introduction

Stochastic processes where small fragments merge and form larger ones are quite useful tools to model physical phenomena at scales ranging from molecular [20] to astronomical ones [24]. The majority of the mathematical literature on such coagulation processes treats mean field models: The rate at which the fragments (clusters) merge is governed only by their sizes - neither the physical location nor their shape affect this rate. See [5] for a review. Stockmayer [20], introduced a mean field model for polymerization where small clusters (sol) merge, however, as soon as a large cluster (gel) forms, it stops growing. In contrast to the mean field models, we consider a model which takes the geometry of the space and the shape of the clusters into account. Following van den Berg, de Lima and Nolin [21], and Aldous [2], we introduce the following adaptation of Stockmayer's model. Let $G = (V, E)$ be a graph which represents the underlying geometry and $N \in \mathbb{N}$. For every vertex $v \in V$, independently from each other, we assign a random time τ_v which is uniformly distributed on $[0, 1]$. At time $t = 0$, all of the vertices of G are closed. As time increases, a vertex v tries to become open at time $t = \tau_v$. It succeeds if and only if all of its neighbours' open clusters (open connected components) at time t have size less than N. Note that as soon as the diameter of a cluster reaches N, it stops growing, i.e freezes. Hence the name N-parameter frozen percolation. Note that we can also consider an edge (bond) version of the model above where edges turn open from closed. This edge version of the process was introduced by van den Berg et al [21].

We are particularly interested in the N-parameter frozen percolation models for large N on graphs such as d dimensional lattices, since they are discrete approximations of the space $\mathbb{R}^d$. Herein we restrict to the case where $d = 2$. We will mainly work on the triangular lattice. We will see that the behaviour of this model rich and interesting too, but in a very different way from the model studied by Aldous [2].

Let us turn to the model introduced and constructed by Aldous [2]. It is the edge version of the model on the binary tree where we replace the parameter N by ∞ in the description above. An edge e of the binary tree opens at time τ_e as long as the open clusters of the endpoints of e are finite. In view of this model, one could also try to construct a similar, so called ∞-parameter, model on the triangular lattice. However Benjamini and Schramm [4] showed that it is impossible. Exactly this non-existence result motivated van den Berg, de Lima and Nolin [21] to extend the model of Aldous for finite parameter N: in this case, the N-parameter frozen percolation process (both the vertex and the edge version) is a finite range interacting particle system, hence the general theory [15] gives existence. One could ask if the N-parameter processes for large but finite N provide a reason for the existence of the ∞-parameter frozen bond percolation and the binary tree and the non-existence of the ∞-parameter frozen site percolation on the triangular lattice. Before we answer this question, let us specify the two dimensional model which plays a central role in this paper.

We will work on the triangular lattice $\mathbb{T} = (V, E)$ with its usual embedding in the plane $\mathbb{R}^2$. That is, the vertex set V is the lattice generated by the vectors $\underline{e}_1 = (1, 0)$ and $\underline{e}_2 = (\cos(\pi/3), \sin(\pi/3))$:

$$V := \{a\underline{e}_1 + b\underline{e}_2 \,|\, a, b \in \mathbb{Z}\}. \tag{1.1}$$

The vertices u and v are neighbours, i.e $(u, v) \in E$ or $u \sim v$ if their L^2 distance is 1. Moreover, we freeze clusters as soon as they reach L^∞ diameter (inherited from $\mathbb{R}^2$) at least N. For the case where the underlying lattice is $\mathbb{Z}^2$ and for different choices for diameters of clusters see the discussion below Conjecture 1.7.

Van den Berg, Kiss and Nolin [23] investigated the edge version of the N-parameter process on the binary tree. They found that as $N \to \infty$, the N-parameter process on the binary tree converges to the ∞-parameter process in some weak sense. This result raises the question if there is a limit of the N-parameter frozen percolation processes on the triangular lattice as N goes to infinity. The non-existence of the ∞-parameter process suggests that the N-parameter model may have a remarkable (anomalous) behaviour in the limit $N \to \infty$. It turns out that there is a limiting process, but this process is, in some sense, trivial:

Theorem 1.1. *As $N \to \infty$ the probability that in the N-parameter frozen percolation process the open cluster of the origin freezes goes to* 0.

To get some intuition for the behaviour of the process, let us for the moment forget about freezing, and call the resulting process the percolation process. That is, at time τ_v the vertex v becomes open no matter how big are the open clusters of its neighbours. Thus at time t, a vertex v is open with probability t independently from the other vertices. Hence at time t we see ordinary site percolation with parameter t. Recall from [18] that the critical parameter for site percolation on the triangular lattice is $p_c = 1/2$. So at each time $t \leq 1/2$ there is no open infinite cluster, and there is a unique infinite open cluster when $t > 1/2$. Moreover, by [1] at time $t < 1/2$, the distribution of the size of the open clusters has an exponential decay. Note that if a site is open in the N-parameter frozen percolation process at time t, then it is also open in the percolation process at time t. Hence at time $t < 1/2$ the

N-parameter frozen percolation process and the percolation process does not differ too much when N is large: even without freezing, for all $K > 0$ the probability that there is an open cluster with diameter at least N in a box with side length KN goes to 0 as $N \to \infty$. To our knowledge, there is no simple argument showing that, roughly speaking, freezing does not take place at times that are essentially bigger than $1/2$, which is one of our main results:

Theorem 1.2. *For all $K > 0$ and $t > 1/2$, the probability that after time t a frozen cluster forms which intersects a given box with side length KN goes to 0 as $N \to \infty$.*

Compare Theorem 1.2 with [2, 21] where it was shown that clusters freeze throughout the time horizon $[1/2, 1]$ for $N \in \mathbb{N} \cup \{\infty\}$ in the edge version of the N-parameter frozen percolation process on the binary tree. (Note that the critical parameter is $1/2$ for site percolation on the binary tree.) As it turns out, our method provides a much stronger result than Theorem 1.2. To state it we need some more notation.

Let $\mathbb{P}$ denote the probability measure corresponding to the percolation process. For a fixed $p \in [0,1]$, we call a vertex $v \in V$ p-open (p-closed), if its τ value is less (greater) than p. We denote by $\mathbb{P}_p$ the distribution of p-open vertices.

We borrow some of the notation from [16]. Recall the definition of V from (1.1). The L^∞ distance of vertices in $\mathbb{T}$ is the L^∞ distance inherited from $\mathbb{R}^2$. That is, for $v, w \in V$ the distance $d(v,w)$ between $v = (v_1, v_2)$ and $w = (w_1, w_2)$ is

$$\begin{aligned} d(v,w) &= \|v - w\|_\infty \\ &= \max\{|v_1 - w_1|, |v_2 - w_2|\}. \end{aligned}$$

For $a, b, c, d \in \mathbb{R}$, with $a < b, c < d$ we define the parallelogram

$$[a,b] \boxtimes [c,d] := \{k\underline{e}_1 + l\underline{e}_2 \,|\, k \in [a,b] \cap \mathbb{Z},\, l \in [c,d] \cap \mathbb{Z}\}.$$

We denote the outer boundary of a set of vertices $S \subseteq V$ by

$$\partial S := \{v \in V \setminus S \,|\, \exists u \in S:\ u \sim v\}.$$

Let $cl(S) = S \cup \partial S$ denote the closure of S. For the parallelogram centred around the vertex v with radius $a > 0$ we write

$$B(v;a) := [-a,a] \boxtimes [-a,a] + v.$$

We denote the annulus centred around $v \in V$ with inner radius $a > 0$ and outer radius $b > a$ by

$$A(v;a,b) := B(v;b) \backslash B(v;a).$$

We call $B(v;a)$ the inner, $B(v;b)$ the outer parallelogram of $A(v;a,b)$.

We say that there is an open (closed) arm in an annulus $A(v;a,b)$ if there is an open (closed) path from $\partial B(v;a)$ to $\partial B(v;b)$ in $A(v;a,b)$. We write o for open and c for closed. A colour sequence of length k is an element of $\{o,c\}^k$. For $\sigma \in \{o,c\}^k$, we denote by $\mathcal{A}_{k,\sigma}(v;a,b)$ the event that there are k disjoint arms in $A(v;a,b)$ such that the vertices of each of the arms are either all open or all closed, moreover, if we take a counter-clockwise ordering of these arms, then their colours follow a cyclic permutation of σ.

In the case where $v = \underline{0} = (0,0)$ we omit the first argument in our notation, that is $B(a) = B(\underline{0};a)$ etc. For the critical arm probabilities we use the notation

$$\pi_{k,\sigma}(a,b) := \mathbb{P}_{1/2}(\mathcal{A}_{k,\sigma}(a,b)). \tag{1.2}$$

In the following we use the near critical parameter scale which was introduced in [12]. For a positive parameter N and $\lambda \in \mathbb{R}$ it is defined as

$$p_\lambda(N) := \frac{1}{2} + \lambda \frac{N^{-2}}{\pi_{4,alt}(1,N)} \tag{1.3}$$

where alt denotes the colour sequence (o,c,o,c).

Before we proceed, let us stop here and let us briefly explain the formula (1.3). Suppose that a vertex v is a closed pivotal vertex, i.e. it is on the boundary of two different open cluster with diameter at least N. The two open clusters provide two disjoint open arms starting from neighbouring vertices of v. Since the open clusters are different, they have to be separated by closed paths, which provide two disjoint closed arms starting from v. Hence the event $\mathcal{A}_{4,alt}(v;1,N)$ occurs. By (1.2), we get that the expected number of pivotal vertices in $B(N)$

is $O\left(N^2\pi_{4,alt}(1,N)\right)$. Let $\lambda > 0$. Let us look at the percolation process in the parallelogram $B(N)$ in the time interval $[1/2, p_\lambda(N)]$. The probability that a vertex opens in this time interval is $p_\lambda(N) - 1/2$. By a combination of (1.2) and (1.3) we see that the expected number of pivotal vertices which open in this interval is $O(1)$. Hence the parameter scale in (1.3) corresponds to the time scale where open clusters of diameter $O(N)$ merge. See [12, 11] for more details.

The considerations above suggest that the parameter scale (1.3) is indeed useful for investigating the N-parameter frozen percolation process. We write $\mathbb{P}_N$ for the probability measure corresponding to the N-parameter frozen percolation process. The following stronger version of Theorem 1.2 is our main result.

Theorem 1.3. *For any $\varepsilon, K > 0$ there exists $\lambda = \lambda(\varepsilon, K)$ and $N_0 = N_0(\varepsilon, K)$ such that*

$$\mathbb{P}_N\left(\textit{a cluster intersecting } B(KN) \textit{ freezes after time } p_\lambda(N)\right) < \varepsilon$$

for all $N \geq N_0$.

In [21] the authors investigated the diameter of the open cluster of the origin at time 1. Their main result is the following.

Definition 1.4. For $t \in [0,1]$ let $C(v;t)$ denote the open cluster of $v \in V$ at time $t \in [0,1]$. We set $C(t) := C(\underline{0};t)$.

Theorem 1.5 (Theorem 1.1 of [21])**.** *For the bond version of the N-parameter frozen percolation on the square lattice we have*

$$\liminf_{N\to\infty} \mathbb{P}_N\left(diam\left(C(1)\right) \in (aN, bN)\right) > 0$$

for $a, b \in (0,1)$ with $a < b$.

Analogous result holds for the (site version of) N-parameter process on the triangular lattice. In the following corollary we supplement this result. It is an extension of Theorem 1.1.

Corollary 1.6. *For any $\varepsilon > 0$ there exists $a = a(\varepsilon), b = b(\varepsilon) \in (0,1)$ with $a < b$ and $N_0 = N_0(\varepsilon)$ such that*

$$\mathbb{P}_N\left(diam\left(C(1) \in (aN, bN)\right)\right) > 1 - \varepsilon$$

for all $N \geq N_0$.

The results above suggest the following intuitive and informal description of the behaviour of N-parameter frozen percolation processes on the triangular lattice for large N: At time 0 all the vertices are open. Then they open independently from each other as in the percolation process till time close to 1/2. Then in the scaling window, frozen clusters form, and by the end of the window, they give a tiling of $\mathbb{T}$ such that all the holes (non-frozen connected components) have diameter less than N but, typically, of order N. After the window, the closed vertices in these holes open as in the percolation process restricted to these holes. At time 1 the non-frozen vertices are all open.

Hence the interesting time scale is (1.3), moreover it raises the question if there is some kind of limiting process which governs the behaviour of the N-parameter frozen percolation processes as $N \to \infty$ in the scaling window (1.3). We have the following conjecture, somewhat informal, conjecture:

Conjecture 1.7. *When we scale space by N and time according to (1.3), we get a non-trivial scaling limit, which is measurable with respect to the near critical ensemble of [12, 11]. Moreover, the scaling limit completely describes the frozen clusters of the N-parameter frozen percolation as $N \to \infty$.*

Let us mention some generalizations of our results. We considered the site version of the N-parameter frozen percolation on the triangular lattice above. Straightforward adaptations of the proofs give the same results for the bond version of the N-parameter frozen percolation on the square lattice. See Remark 3.7 for more details. Our results remain valid when use some different distance instead of the L^∞ distance in the definition of the N-parameter frozen percolation process, as long as the used distance resembles the L^∞ distance. Examples of such distances include the L^p distances for some $p \geq 1$, or when we rotate the lattice $\mathbb{T}$. Finally let us mention that when we freeze clusters when their *volume* (number of its vertices) reach N, then we get a quite different process.

Let us briefly discuss some related results. A version of the N-parameter frozen percolation process on $\mathbb{Z}$ and the binary tree was investigated in [8]. We already considered [2] where Aldous introduced the ∞-parameter frozen percolation process on the binary tree. However, we did not mention that this model has an other interesting, so called self organized critical (SOC), behaviour: For all $t > 1/2$, the distribution of the active clusters at time t have the same distribution as critical clusters. Clearly, the N-parameter frozen percolation process on the triangular lattice does not have this property. A mean field version of the frozen percolation model on the complete graph investigated by Ráth in [17]. He showed that this model has similar SOC properties. Finally, we refer to [6] where a dynamics similar to frozen percolation was investigated on uniform Cayley trees.

The organization of the paper is the following. In Section 2, we introduce some more notation, and briefly discuss the results from percolation theory required to prove our main result: We start with some classical correlation inequalities in Section 2.1. In Section 2.2 we introduce mixed arm events where some of the arms can use only the upper half of the annulus, while others can use the whole annulus. Here we also recall some of their well-known properties and discuss some new ones. In particular, we note that the exponent of the arm events increases when we increase the number of arms which have to stay in the upper half plane. The proof of this statement is postponed to Section A.1 of the Appendix. In Section 2.3 we describe the connection between the correlation length with the near critical scaling (1.3). We prove Theorem 1.3 and Corollary 1.6 in Section 3 assuming two technical results Proposition 3.5 and 3.6. In Section 4 we introduce some more notation and the notion of thick paths. There we prove Proposition 3.6. In this proof a deterministic (combinatorial/geometric) result, Lemma 4.5, plays an important role. The proof of this lemma is postponed to Section A.2 of the Appendix. The most technical part of the paper is Section 5 where we prove Proposition 5. In Section 5.1 and 5.2 we investigate the vertical position of the lowest point of the lowest closed crossing in regions with half open half closed boundary conditions. We combine these results with the ones in Section 2 and conclude the proof of Proposition 3.5 in Section 4. This finishes the proof of the main result.

Acknowledgement

The author thanks Jacob van den Berg, Federico Camia, Pierre Nolin and Gábor Pete for fruitful discussions. He thanks Gábor Pete and Artëm Sapozhnikov for calling his attention to Lemma 2.11, which was instrumental in the proof of the main result. He is grateful to Jacob van den Berg for his numerous comments on the earlier versions of the paper.

2 Preliminary results on near critical percolation

We recall some classical results from percolation theory in this section. With suitable modifications, the results of this section also hold for bond percolation on the square lattice unless it is indicated otherwise.

2.1 Correlation inequalities

We use the following two inequalities throughout the paper. See Section 2.2 and 2.3 of [13] for more details. We refer to the first theorem as BK, and as FKG for the second.

Theorem 2.1 (BK). *Let A, B be arbitrary events, then*

$$\mathbb{P}_p\left(A \Box B\right) \leq \mathbb{P}_p\left(A\right)\mathbb{P}_p\left(B\right).$$

Definition 2.2. Let $A \subset \{o,c\}^V$ and $U \subseteq V$. We say that an event $A \subset \{o,c\}^V$ is increasing (decreasing) in the configuration in U, if for all $\omega \in A$ we have $\omega' \in A$ where

$$\omega'(v) = \begin{cases} \omega(v) \text{ or } o\ (c) & \text{for } v \in U \\ \omega(v) & \text{for } v \in V \setminus U. \end{cases}$$

That is, turning some closed (open) vertices in U into open (closed) ones can only help the occurrence of A. In the case where $U = V$ we simply say that A is an increasing (decreasing) event.

Theorem 2.3 (FKG). *For any pair of increasing events A, B we have*

$$\mathbb{P}_p\left(A \cap B\right) \geq \mathbb{P}_p\left(A\right)\mathbb{P}_p\left(B\right).$$

2.2 Mixed arm events, critical arm exponents

Recall the definition of arm events from the introduction. There the arms were allowed to use the whole annulus. We introduce the mixed arm events, where some of the arms lie in the upper half of the annulus, while others can use the whole annulus:

Definition 2.4. Let $l, k \in \mathbb{N}$ with $0 \leq l \leq k$, and a colour sequence σ of length k. Let $v \in V$ and $a, b \in (1, \infty)$ with $a < b$. The the full plane k, l mixed arm event with colour sequence σ in the annulus $A(v; a, b)$ is denoted by $\mathcal{A}_{k,l,\sigma}(v; a, b)$. It is the normal k arm event $\mathcal{A}_{k,\sigma}(v; a, b)$ of the Introduction with the extra the condition that there is a counter-clockwise ordering of the arms such that the colour of the arms follow σ, and the first l arms lie in the half annulus $A(v; a, b) \cap (\mathbb{Z} \boxtimes [0, \infty) + v)$. When $v = 0$, we omit the first argument from these notations.

We extend the definition (1.2) for mixed arm events by defining

$$\pi_{k,l,\sigma}(a, b) := \mathbb{P}_{1/2}(\mathcal{A}_{k,l,\sigma}(a, b)).$$

Remark 2.5. In the case $k = l$, we get the so called half plane arm events.

We fix $n_0(k) = 10k$ for $k \in \mathbb{N}$. Note that the event $\mathcal{A}_{k,l,\sigma}(n, N)$ is non-empty whenever $n_0(k) < n < N$. Let us summarize the known critical arm exponents for site percolation on the triangular lattice. To our knowledge, Theorem 2.6 in its generality is not known to hold for bond percolation on $\mathbb{Z}^2$.

Theorem 2.6 (Theorem 3 and 4 of [19]). *Let $l, k \in \mathbb{N}$ and σ be a colour sequence of length k. We define $a_{k,l}(\sigma)$*

- *for $k = 1, l = 0$ and any colour sequence σ as*

$$\alpha_{1,0}(\sigma) := \frac{5}{48},$$

- *for $k > 1$ and $l = 0$, when σ contains both colours, as*

$$\alpha_{k,0}(\sigma) := \frac{k^2 - 1}{12},$$

- *for $k = l \geq 1$ and any colour sequence σ as*

$$\alpha_{k,k}(\sigma) := \frac{k(k+1)}{6}.$$

In these cases we have

$$\pi_{k,l,\sigma}(n_0(k), N) = N^{-\alpha_{k,l}(\sigma) + o(1)}$$

as $N \to \infty$,

To our knowledge, for general k and l, neither the value, nor the existence of the exponents is not known. We expect that the exponents do exist. We will see in Proposition A.5, that if $\alpha_{k,l}(\sigma)$ and $\alpha_{k,m}(\sigma)$ exists for some $k, l, m \in \mathbb{N}$ and $\sigma \in \{o, c\}^k$ with $m < l$, then $\alpha_{k,m}(\sigma) < \alpha_{k,l}$. Since we do not need such general result, we only prove the following proposition in detail.

Proposition 2.7. *For any $k \geq 1$, there are positive constants $c = c(k)$, $\varepsilon = \varepsilon(k)$ such that*

$$\pi_{k,l,\sigma}(n_0(k), N) \leq cN^{-\varepsilon} \pi_{k,0,\sigma}(n_0(k), N) \tag{2.1}$$

for $l = 1, 2, \ldots, k$ uniformly in N and in the colour sequence σ.

Remark 2.8. (i) We do not need the exact values of the critical exponents of Theorem 2.6. For our purposes it is enough to show that certain arm events have exponents at least 2.

(ii) Proposition 2.7 and its generalization also hold for mixed arm events in bond percolation on the square lattice.

Proof of Proposition 2.7. Proposition 2.7 is a simple corollary of Proposition A.3 of the Appendix. Loosely speaking, it states that conditioning on the event that we have k arms in $A(a, b)$, these arms wind around the centre of the annulus $O(\log(b/a))$ many times with probability $1 - \left(\frac{b}{a}\right)^{\kappa}$ for some $\kappa > 0$. The proof of Proposition A.3 can be found in the Appendix. ☐

Remark 2.9. Recall that we do not know in general if the exponents $\alpha_{k,l}(\sigma)$ exist or not. Nonetheless, on the triangular lattice, Proposition 2.7 and Theorem 2.6 and the BK inequality (Theorem 2.1) give that for any colour sequence σ, there is an upper bound with exponent strictly larger than 2 for $\pi_{k,l,\sigma}(n_0(k), N)$ when

- $k \geq 6$, and $l \geq 0$, or
- $k \geq 5$ and $l \geq 1$, or
- $k \geq 4$ and $l \geq 3$.

For arm events with exponents larger than 2 in the case of bond percolation on the square lattice see Remark 2.14 below.

An other well-known attribute of critical arm events is their quasi-multiplicative property. For the full plane, respectively for half plane, arm events this property is shown to hold in Proposition 17 of [16], respectively in Section 1.4.6 of [16]. Simple modifications of these arguments apply to mixed arm events. We introduce the notation $\asymp$ when the ratio of the two quantities is bounded away from 0 and ∞. With this notation we get:

Proposition 2.10. *Let $k \geq 1$ and $\sigma \in \{o, c\}^k$. Then*

$$\pi_{k,l,\sigma}(n_1, n_2)\, \pi_{k,l,\sigma}(n_2, n_3) \asymp \pi_{k,l,\sigma}(n_1, n_3)$$

uniformly in $n_0(k) \leq n_1 \leq n_2 \leq n_3$.

In the following lemma we consider arm events where the open arms are p-open and the closed arms are q-closed where $p, q \in [0,1]$ with p not necessarily equal to q. When p and q are of the form (1.3), then we call these arm events near critical arm events. In this case the probabilities of these events are comparable to critical arm event probabilities. The following lemma is a generalization of Lemma 2.1 of [10] and Lemma 6.3 of [9].

Lemma 2.11. *Let $v \in V$, $\lambda_1, \lambda_2 \in \mathbb{R}$ and $a, b \in (0,1)$ with $a < b$. Let $\mathcal{A}^{\lambda_1,\lambda_2,N}_{k,l,\sigma}(v; aN, bN)$ denote the modification of the event $\mathcal{A}_{k,l,\sigma}(v; aN, bN)$ where the open arms are $p_{\lambda_2}(N)$-open and the closed arms are $p_{\lambda_1}(N)$-closed. Then there are positive constants $c = c(\lambda_1, \lambda_2, k)$ and $N_0 = N_0(\lambda_1, \lambda_2, a, b, k)$ such that*

$$\mathbb{P}\left(\mathcal{A}^{\lambda_1,\lambda_2,N}_{k,l,\sigma}(v; aN, bN)\right) \leq c\pi_{k,l,\sigma}(aN, bN)$$

for $N \geq N_0$.

Proof of Lemma 2.11. It follows from either of the proof of Lemma 2.1 of [10] or from the adaptation of the proof of Lemma 6.3 of [9] to the triangular lattice. ☐

In the following events we collect some of the near critical arm events which have upper bounds with exponents strictly larger than 2. These events play a crucial role in our main result.

Definition 2.12. Let $a, b \in (0,1)$, $\lambda_1, \lambda_2 \in \mathbb{R}$, $K > 0$ and $N \in \mathbb{N}$ with $a < b$. Let $\mathcal{NA}^c(a, b, \lambda_1, \lambda_2, K, N)$ denote the union of the events $\mathcal{A}^{\lambda_1,\lambda_2,N}_{k,l,\sigma}(v; aN, bN)$ for $(k,l) \in \{(4,3), (5,1), (6,0)\}$, $\sigma \in \{o,c\}^k$ as well as the versions of these events where the half plane arms can only use the lower, left or right half of the annulus $A(v; aN, bN)$. We define $\mathcal{NA}(a, b, \lambda_1, \lambda_2, K, N)$ as the complement of the event above.

We show that that for fixed b, K, λ_1 and λ_2, we can set $a \in (0,1)$ so that the probability of $\mathcal{NA}(a, b, \lambda_1, \lambda_2, K, N)$ becomes as small as we require for large N. More precisely, we prove the following:

Corollary 2.13. *There is $\tilde{\varepsilon} > 0$ such that for all $a, b \in (0,1)$, with $a < b$ and $\lambda_1, \lambda_2 \in \mathbb{R}$ there are positive constants $c = c(\lambda_1, \lambda_2, K)$ and $N_0 = N_0(a, b, \lambda_1, \lambda_2, K)$ such that*

$$\mathbb{P}(\mathcal{NA}(a, b, \lambda_1, \lambda_2, K, N)) \geq 1 - c\frac{a^{\tilde{\varepsilon}}}{b^{2+\tilde{\varepsilon}}}$$

for $N \geq N_0$.

Proof of Corollary 2.13. Suppose that one of the arm events in Definition 2.12, for example $\mathcal{A}^{\lambda_1,\lambda_2,N}_{k,l,\sigma}(v; aN, bN)$ for some $v \in B(KN)$, occurs. Then the event $\mathcal{A}^{\lambda_1,\lambda_2,N}_{k,l,\sigma}\left(\lfloor 2aN \rfloor z; 2aN, \frac{b}{2}N\right)$ occurs for some $z \in V$ with $z \in B\left(\lceil \frac{a+K}{2a} \rceil\right)$.

Combination of Remark 2.9 and Lemma 2.11 gives that there are constants $c' = c'(\lambda_1, \lambda_2)$ and $N_0 = N_0(a, b, \lambda_1, \lambda_2)$, and a universal constant $\tilde{\varepsilon} > 0$ such that the probability of one of these events is at most

$$c' \left(\frac{2a}{b/2} \right)^{2+\varepsilon} \tag{2.2}$$

for $N \geq N_0$. The same argument works for other arm events which appear in Definition 2.12, and provide an upper bound similar to (2.2). Hence (2.2) combined with $\left| B\left(\left\lceil \frac{a+K}{2a} \right\rceil \right) \right| = O\left(a^{-2}\right)$ concludes the proof of Corollary 2.13. $\square$

Remark 2.14. To our knowledge it is not known if the direct analogue of Corollary 2.13 holds on the square lattice. The reason is that the exponent $\alpha_{5,0}(\sigma)$ and $\alpha_{3,3}(\sigma)$ is not known for general σ. See Remark 26 of [16].

We recall the proof of Theorem 24 and Remark 26 of [16], where it is shown that $\alpha_{5,0}(o, c, o, o, c) = 2$ and $\alpha_{3,3}(c, o, c) = 2$ on the square lattice. This implies that a version of Corollary 2.13 holds for the square lattice if we modify Definition 2.12 so that we only forbid the occurrence of those arm events where the required set of arms contain

- three half plane arm events with colour sequence (o, c, o) or (c, o, c), or
- five full plane arms with colour sequence (o, c, o, o, c) or (c, o, c, c, o)

as a subset.

2.3 Near-critical scaling and correlation length

Recall that in Section 1 we already gave an explanation for the near critical parameter scale (1.3). In this section we give an other interpretation of this parameter scale, which is connected to the correlation length introduced by Kesten in [14].

We say that there is an open (closed) horizontal (vertical) crossing of a parallelogram $B := [a, b] \boxtimes [c, d]$ if there is an open (closed) path connecting $\{\lceil a \rceil\} \boxtimes [c, d]$ and $\{\lfloor b \rfloor\} \boxtimes [c, d]$ in$[a, b] \boxtimes [c, d]$. For the event that there is an open (closed) horizontal crossing of B we use the notation $\mathcal{H}_o(B)$ ($\mathcal{H}_c(B)$). One can define similar events for vertical crossings, which we denote by by $\mathcal{V}_o(B)$ and $\mathcal{V}_c(B)$. For $\varepsilon \in (0, 1/2)$ the correlation length is defined as

$$L_\varepsilon(p) = \begin{cases} \min\left\{ n \,|\, \mathbb{P}_p\left(\mathcal{H}_o(B(n))\right) \leq \varepsilon \right\} & \text{when } p < p_c \\ \min\left\{ n \,|\, \mathbb{P}_p\left(\mathcal{H}_o(B(n))\right) \geq 1 - \varepsilon \right\} & \text{when } p > p_c. \end{cases}$$

Remark 2.15. The particular choice of ε is not important in this definition. Indeed, Corollary 37 of [16], or alternatively Corollary 2 of [14], gives that

$$L_\varepsilon(p) \asymp L_{\varepsilon'}(p)$$

for any $\varepsilon, \varepsilon' \in (0, 1/2)$ uniformly in $p \in (0, 1)$.

We show that the control over the near critical parameter λ gives a control over the correlation length in Corollary 2.17 and 2.18 below. Recall the remark after Lemma 8 of [14]:

Proposition 2.16. *For any fixed* $\varepsilon \in (0, 1/2)$, *we have*

$$|p - p_c| \left(L_\varepsilon(p)\right)^2 \pi_{4,0,alt}\left(1, L_\varepsilon(p)\right) \asymp 1$$

uniformly for $p \neq 1/2$.

Note that for fixed $\varepsilon > 0$, the correlation length $L_\varepsilon(p)$ is a decreasing (increasing) function of p for $p > p_c$ ($p < p_c$). Combination of this and Proposition 2.10 we get:

Corollary 2.17. *For all* $\lambda \in \mathbb{R} \setminus \{0\}$ *and* $\varepsilon \in (0, 1/2)$,

$$L_\varepsilon\left(p_\lambda(N)\right) \asymp N. \tag{2.3}$$

Corollary 2.18. *For any* $C>0$ *and* $\varepsilon\in(0,1/2)$ *there exits* $\lambda_1=\lambda_1(C,\varepsilon)>0$ *and* $N_1=N_1(C,\varepsilon)$ *such that for any* $\lambda\in\mathbb{R}$ *with* $|\lambda|\geq\lambda_1$ *we have*

$$L_\varepsilon(p_\lambda(N))\leq CN$$

for $N\geq N_1$*. Also, for any* $c>0$*, and* $\varepsilon\in(0,1/2)$ *there exists* $\lambda_2(c,\varepsilon)>0$ *and* $N_2=N_2(c,\varepsilon)$ *such that for any* $\lambda\in\mathbb{R}\setminus\{0\}$ *with* $|\lambda|\leq\lambda_2$ *we have*

$$L_\varepsilon(p_\lambda(N))\geq cN$$

for $N\geq N_2$*.*

Remark 2.19. On the triangular lattice, a ratio limit theorem for $\pi_{4,0,alt}$, Proposition 4.7 of [12] holds. This combined with the definition of $L_\varepsilon(p)$, and Proposition 2.16 shows that the following stronger statement holds on the triangular lattice:

Claim. For all $\lambda_1,\lambda_2\in\mathbb{R}$ with $\lambda_1\leq\lambda_2$, $\lambda_1\lambda_2>0$ and $\varepsilon\in(0,1/2)$ there are positive constants $c=c(\varepsilon),C=C(\varepsilon)$ and $N_0=N_0(\varepsilon,\lambda_1,\lambda_2)$ such that

$$cN|\lambda|^{-4/3}\leq L_\varepsilon(p_\lambda(N))\leq CN|\lambda|^{-4/3}$$

for all $\lambda\in[\lambda_1,\lambda_2]$ and $N\geq N_0$.

Standard Russo-Seymour-Welsh (RSW) techniques and the definition of the correlation length give that the control over the correlation length gives a control over the crossing probabilities of parallelograms. This combined with the two corollaries above show that that the control over the near critical parameter gives a control over the crossing probabilities. See Corollary 2.20 and 2.21 below:

Corollary 2.20. *For all* $\lambda\in\mathbb{R}$ *and* $a,b\in(0,\infty)$*, there are constants* $c=c(a,b,\lambda)\in(0,1)$*,* $C=C(a,b,\lambda)\in(0,1)$ *and* $N_0=N_0(a,b,\lambda)$ *with such that*

$$\begin{gathered}c<\mathbb{P}_{p_\lambda(N)}(\mathcal{H}_o([0,aN]\boxtimes[0,bN]))<C\\ c<\mathbb{P}_{p_\lambda(N)}(\mathcal{H}_c([0,aN]\boxtimes[0,bN]))<C\end{gathered}$$

for $N\geq N_0$*.*

Corollary 2.21. *Let* $\delta\in(0,1)$*, and* $a,b\in(0,\infty)$*. There exists* $\lambda_1=\lambda_1(\delta,a,b)>0$ *and* $N_1=N_1(\delta,a,b)$ *such that for all* $\lambda\geq\lambda_1$

$$\mathbb{P}_{p_\lambda(N)}(\mathcal{H}_o([0,aN]\boxtimes[0,bN]))>1-\delta$$

for $N\geq N_1$*. Furthermore, there exists* $\lambda_2=\lambda_2(\delta,a,b)<0$ *and* $N_2=N_2(\delta,a,b)$ *such that for all* $\lambda\leq\lambda_2$

$$\mathbb{P}_{p_\lambda(N)}(\mathcal{H}_c([0,aN]\boxtimes[0,bN]))>1-\delta$$

for $N\geq N_2$*.*

Similar RSW techniques show that it is unlikely to have crossing in a thin and long parallelogram in the hard direction in the critical window. See Remark 40 [16] for more details.

Corollary 2.22. *Let* $\lambda\in\mathbb{R}$*, and* $a,b\in(0,1)$*. There exists positive constants* $c=c(\lambda),C=C(\lambda)$ *and* $N_0=N_0(\lambda,a,b)$ *such that*

$$\mathbb{P}_{p_\lambda(N)}(\mathcal{H}_o([0,aN]\boxtimes[0,bN]))\leq C\exp\left(-c\frac{a}{b}\right)$$

for $N\geq N_0$*.*

The following event plays a crucial role in the proof of our main result.

Definition 2.23. Let $a,b\in(0,1)$, $\lambda_1,\lambda_2\in\mathbb{R}$, and $N\in\mathbb{N}$ with $a<b$. Let $\mathcal{NC}(a,b,\lambda_1,\lambda_2,K,N)$ denote the event that for all parallelograms $B=[0,aN]\boxtimes[0,bN]+z$ with $z\in B(KN)$, there is neither a $p_{\lambda_1}(N)$-open nor a $p_{\lambda_2}(N)$-closed horizontal crossing in B.

The following Corollary 2.24 follows from Corollary 2.22 by arguments analogous to the proof of Corollary 2.13.

Corollary 2.24. *Let $a, b \in (0,1)$, $\lambda_1, \lambda_2 \in \mathbb{R}$, and $N \in \mathbb{N}$ with $a < b$. There are positive constants $c = c(\lambda_1, \lambda_2)$, $C = C(\lambda_1, \lambda_2)$ and $N_0 = N_0(a, b, \lambda_1, \lambda_2)$ such that*

$$\mathbb{P}(\mathcal{NC}(a, b, \lambda_1, \lambda_2, K, N)) \geq 1 - Ca^{-2} \exp\left(-c\frac{b}{a}\right)$$

for $N \geq N_0$.

We finish this section by stating two lemmas which will be used explicitly in the proof of our main result.

Lemma 2.25. *For any fixed $\lambda \in \mathbb{R}$, for any $a, b \in (0, \infty)$ and $\varepsilon > 0$, there is are positive integer $K = K(\lambda, a, b, \varepsilon)$ and $N_0 = N_0(\lambda, a, b, \varepsilon)$ such that*

$$\mathbb{P}_{p_\lambda(N)}(\textit{there are at least } K \textit{ disjoint closed arms in } A(aN, bN)) < \varepsilon$$

for $N \geq N_0$.

Proof of Lemma 2.25. This is a consequence of Corollary 2.20 and the BK inequality (Theorem 2.1). The proof also also appears in the proof of Lemma 15 of [16]. □

Definition 2.26. Let $a, b, c, d, f \in \mathbb{R}$ with $a \leq b$, $c \leq d$ and $f > 0$. We say that there is an open (closed) f-net in $B = [a, b] \boxtimes [c, d]$ if there is an open (closed) vertical crossing in the parallelograms $[a + i\lfloor f \rfloor, a + (i+1)\lfloor f \rfloor - 1] \boxtimes [c, d]$, and there is an open (closed) horizontal crossing in the parallelograms $[a, b] \boxtimes [c + j\lfloor f \rfloor, c + (j+1)\lfloor f \rfloor - 1]$ for $i = 0, 1, \ldots, \lfloor (b-a) / \lfloor f \rfloor \rfloor$ and $j = 0, 1, \ldots, \lfloor (d-c) / \lfloor f \rfloor \rfloor$.

For $\lambda \in \mathbb{R}$ and $\delta \in (0, \infty)$, $\mathcal{N}_c(\lambda, \delta, K, N)$ ($\mathcal{N}_o(\lambda, \delta, K, N)$) denotes the event that there is a $p_\lambda(N)$-closed ($p_\lambda(N)$-open) δN-net in $B(KN)$.

Lemma 2.27. *Let $\varepsilon, \delta, K > 0$. There exists $\lambda_1 = \lambda_1(\varepsilon, \delta, K) \in \mathbb{R}$ and $N_1 = N_1(\varepsilon, \delta, K)$ such that*

$$\mathbb{P}(\mathcal{N}_o(\lambda_1, \delta, K, N)) > 1 - \varepsilon$$

for $N \geq N_1$. Moreover there exists $\lambda_2 = \lambda_2(\varepsilon, \delta, K) \in \mathbb{R}$ and $N_2 = N_2(\varepsilon, \delta, K)$ such that

$$\mathbb{P}(\mathcal{N}_c(\lambda_2, \delta, K, N)) > 1 - \varepsilon$$

for $N \geq N_2$.

Proof of Lemma 2.27. This is a consequence of Corollary 2.21 and Theorem 2.3. □

3 Proof of the main results

We prove our main results Theorem 1.3 and Corollary 1.6 in this section assuming Proposition 3.5 and 3.6.

Definition 3.1. In the N-parameter frozen percolation process we call a vertex *frozen* at some time $t \in [0,1]$, if either it or one of its neighbours have an open cluster with diameter bigger than N at time t. If a site is not frozen at time t, then we say it is *active* at time t. Note that both frozen and active sites can be open or closed. We say that F is a *(open) frozen cluster* at time $t \in [0,1]$ if it is a connected component of the open vertices at time t with $diam(F) \geq N$. In the case where $t = 1$, we simply say that F is a frozen cluster.

Recall Definition 2.26. We observe the following.

Observation 3.2. *Let $K > 0$ and $N \in \mathbb{N}$. Then in the N-parameter frozen percolation process there is no frozen cluster at time $p_\lambda(N)$ in $B(KN)$ on the event $\mathcal{N}_c(\lambda, 1/6, K+2, N)$. Hence on $\mathcal{N}_c(\lambda, 1/6, K+2, N)$, a vertex in $B(KN)$ is open (closed) in the N-parameter frozen percolation process at time $p_\lambda(N)$ if and only if it is $p_\lambda(N)$-open ($p_\lambda(N)$-closed).*

We show that the number of frozen clusters intersecting $B(KN)$ in the N-parameter frozen percolation process is tight in N.

Lemma 3.3. *Let $K > 0$ and $N \in \mathbb{N}$. Let $FC(t) = FC(t, K, N)$ denote the number of frozen clusters intersecting $B(KN)$ at time $t \in [0,1]$ in the N-parameter frozen percolation process. Then for all $\varepsilon > 0$ there exists $L = L(\varepsilon, K)$ and $N_0 = N_0(\varepsilon, K)$ such that*

$$\mathbb{P}_N(FC(1) > L) < \varepsilon$$

for $N \geq N_0$.

Proof of Lemma 3.3. By Lemma 2.27 we set $\lambda = \lambda(\varepsilon, K) \in \mathbb{R}$ such that

$$\mathbb{P}_N\left(\mathcal{N}_c\left(\lambda, 1/6, K+4, N\right)\right) > 1 - \frac{1}{2}\varepsilon \tag{3.1}$$

for $N \geq N_1(\varepsilon, K)$. Let F be an open frozen cluster which intersects $B(KN)$. From Observation 3.2 we get the vertices of ∂F are closed at $p_\lambda(N)$ in the N-parameter percolation process on the event $\mathcal{N}_c(\lambda, 1/6, K+4, N)$.

Let us cover the parallelogram $B(KN)$ with the annuli

$$A_z = A\left(\lfloor N/20 \rfloor z; \lfloor N/20 \rfloor, \lfloor N/10 \rfloor\right) \text{ with } z \in B\left(\lceil 20K \rceil\right).$$

Suppose that there is an open frozen cluster in the N-parameter frozen percolation which has a vertex in $B(KN)$. The construction of the annuli above gives that there is $z \in B(\lceil 20K \rceil)$ such that $B(\lfloor N/20 \rfloor z; \lfloor N/20 \rfloor)$, the inner parallelogram of A_z, contains a vertex of this open frozen cluster. Since the diameter of $B(\lfloor N/20 \rfloor z; \lfloor N/10 \rfloor)$ is less than N, this cluster has to cross the annulus A_z. Hence for each open frozen cluster intersecting $B(KN)$, we find at least one open frozen crossing of an annulus A_z. Moreover, if there are $k \geq 2$ different frozen clusters crossing some annulus A_z, then there are at least k disjoint closed frozen arms which separate the open frozen clusters in A_z at time 1. By the arguments above, these arms are $p_\lambda(N)$-closed. Thus the number of different frozen clusters intersecting $B(\lfloor N/20 \rfloor z; \lfloor N/20 \rfloor)$ is bounded above by $1 \vee l_z$, where l_z is the number of disjoint $p_\lambda(N)$-closed arms of A_z. Hence by the translation variance of the N-parameter frozen percolation process we have

$$\begin{aligned}
\mathbb{P}_N\left(FC(1) \geq L, \mathcal{N}_c(\lambda, 1/24)\right) &\leq \mathbb{P}_{p_\lambda(N)}\left(\sum_{z \in B(\lceil 20K \rceil)} (1 \vee l_z) \geq L\right) \\
&\leq \mathbb{P}_{p_\lambda(N)}\left(\exists z \in B(\lceil 20K \rceil) \text{ such that } l_z \geq \left(2\lceil 20K \rceil + 1\right)^{-2} L\right) \\
&\leq \left(2\lceil 20K \rceil + 1\right)^2 \mathbb{P}_{p_\lambda(N)}\left(l_0 \geq \left(2\lceil 20K \rceil + 1\right)^{-2} L\right) \qquad (3.2)
\end{aligned}$$

By Lemma 2.25 we set $L = L(\varepsilon, K) \geq \left(2\lceil 20K \rceil + 1\right)^2$ and $N_2 = N_2(\varepsilon, K)$ such that

$$\mathbb{P}_{p_\lambda(N)}\left(l_0 \geq L/200^2\right) < \frac{1}{2}\left(2\lceil 20K \rceil + 1\right)^{-2} \varepsilon$$

for $N \geq N_2$. This combined with (3.2) gives that

$$\mathbb{P}_N\left(FC(1) \geq L, \mathcal{N}_c(\lambda, 1/6, K+4, N)\right) < \frac{1}{2}\varepsilon \tag{3.3}$$

for $N \geq N_2$. We set $N_0 := N_1 \vee N_2$. A combination of (3.1) and (3.3) finishes the proof of Lemma 3.3. ☐

Definition 3.4. For $v \in V$ and $\lambda \in \mathbb{R}$ let $\mathcal{C}_a(v; \lambda) = \mathcal{C}_a(v; \lambda, N)$ denote the active cluster of v in the N-parameter frozen percolation process at time $p_\lambda(N)$. We omit the first argument from the notation above when $v = \underline{0}$.

We state the two propositions below which play a crucial role in the proof of Theorem 1.3. The proof of these propositions are rather technical, so we postpone them to the next section. The first proposition shows that for $\alpha > 0$, it is unlikely to have an active cluster at time $p_\lambda(N)$ which intersects $B(KN)$ and has diameter close to αN.

Proposition 3.5. *For all $\lambda \in \mathbb{R}$ and $\varepsilon, K, \alpha > 0$, there exist $\theta = \theta(\lambda, \alpha, \varepsilon, K) \in (0, 1/2)$ and $N_0 = N_0(\lambda, \alpha, \varepsilon, K)$ such that*

$$\mathbb{P}_N\left(\exists v \in B(KN) \text{ s.t. } diam\left(\mathcal{C}_a(v; \lambda)\right) \in \left((\alpha - \theta)N, (\alpha + \theta)N\right)\right) < \varepsilon$$

for $N \geq N_0$.

The second proposition claims that if there is a vertex v such that $diam\left(\mathcal{C}_a(v; \lambda_1, N)\right) \geq (1+\theta)N$ then some part of $\mathcal{C}_a(v; \lambda_1, N)$ freezes 'soon':

Proposition 3.6. *Let $\theta \in (0,1)$, $\varepsilon > 0$ and $\lambda_1 K, \in \mathbb{R}$. Recall the notation $FC(t, K+2, N)$ from Lemma 3.3. There exists $\lambda_2 = \lambda_2(\lambda_1, \theta, \varepsilon)$ and $N_0 = N_0(\lambda_1, \theta, \varepsilon)$ such that the probability of the intersection of the events*

- *$\exists v \in B(KN)$ such that $diam\left(\mathcal{C}_a(v; \lambda_1, N)\right) \geq (1+\theta)N$, and*

- *none of the clusters intersecting* $B\left(\left(K+2\right)N\right)$ *freeze in the time interval* $\left(p_{\lambda_1}\left(N\right), p_{\lambda_2}\left(N\right)\right]$, *i.e.*

$$FC\left(p_{\lambda_1}\left(N\right), K+2, N\right) = FC\left(p_{\lambda_2}\left(N\right), K+2, N\right)$$

is less than ε *for* $N \geq N_0$.

Before we turn to the proof of our main results we make a remark on how to adapt the proofs for the N-parameter frozen bond percolation process on the square lattice.

Remark 3.7. The arguments in Section 3, 4, 5 and in the Appendix can be easily adapted to the N-parameter frozen bond percolation on the square lattice. Some care is required when we use some Corollary 2.13: As we already noted in Remark 2.14, the direct analogue of Corollary 2.13 does not hold on the square lattice. However, one can check that the version of Corollary 2.13 which was proposed in Remark 2.14 is enough for the proofs appearing in Section 3, 4, 5.

3.1 Proof of Theorem 1.3

Proof of Theorem 1.3. The proof follows the following informal strategy. Consider the following procedure. We set $\lambda_1 = 0$. We look at the N-parameter percolation process at time $p_{\lambda_1}\left(N\right)$. We have two cases.

In the first case all the active clusters at time $p_{\lambda_1}\left(N\right)$ intersecting $B\left(KN\right)$ have diameter less than N. Hence no cluster intersecting $B\left(KN\right)$ can freeze after $p_{\lambda_1}\left(N\right)$. We terminate the procedure.

In the second case there is $v \in B\left(KN\right)$ such that the active cluster $\mathcal{C}_a\left(v; \lambda_1, N\right)$ has diameter at least N. Using Proposition 3.5 we set θ_1 such that the diameter of this cluster is at least $\left(1+\theta_1\right)N$ with probability close to 1. If $diam\left(\mathcal{C}_a\left(v; \lambda_1, N\right)\right) \leq \left(1+\theta_1\right)N$, then we stop the procedure. If $diam\left(\mathcal{C}_a\left(v; \lambda_1, N\right)\right) > \left(1+\theta_1\right)N$, then using Proposition 3.6 we set $\lambda_2 \geq \lambda_1$ such that some part of $\mathcal{C}_a\left(v; \lambda_1, N\right) \cap B\left(\left(K+2\right)N\right)$ freezes in the time interval $\left[p_{\lambda_1}\left(N\right), p_{\lambda_2}\left(N\right)\right]$ with probability close to 1. If indeed some part of $\mathcal{C}_a\left(v; \lambda_1, N\right) \cap B\left(\left(K+2\right)N\right)$ freezes in the time interval $\left[p_{\lambda_1}\left(N\right), p_{\lambda_2}\left(N\right)\right]$, then we iterate the procedure starting from time $p_{\lambda_2}\left(N\right)$. Otherwise we terminate the procedure.

Using Lemma 3.3 we set L such that the event where there are at least L frozen clusters intersecting $B\left(\left(K+2\right)N\right)$ at time 1 has probability smaller than $\varepsilon/2$. In each step of the procedure either the procedure stops, or the the number frozen clusters intersecting $B\left(\left(K+2\right)N\right)$ increases by at least 1. Hence the event that the procedure runs for at least L steps has probability at most $\varepsilon/2$.

Moreover, we set the parameters λ_i, θ_i for $i \geq 1$ above such that with probability at least $1-\varepsilon/2$ we terminate the procedure when there are no active clusters intersecting $B\left(KN\right)$ with diameter at least N. Thus with probability at least $1-\varepsilon$ the procedure stops within L steps, and we stop when there are no active clusters with diameter at least N intersecting $B\left(KN\right)$. Hence $\lambda = \lambda_{L+1}$ satisfies the conditions of Theorem 1.3, which finishes the proof of Theorem 1.3.

Let us turn to the precise proof. By Lemma 3.3, there is $L = L\left(\varepsilon, K\right)$ and $N_1' = N_1'\left(\varepsilon, K\right)$ such that

$$\mathbb{P}_N\left(FC\left(1, K+2, N\right) \geq L\right) \leq \varepsilon/2, \tag{3.4}$$

where $F\left(t, K+2, N\right)$ counts the number of frozen clusters intersecting $B\left(\left(K+2\right)N\right)$ at time $t \in \left[0,1\right]$.

We define the deterministic sequence $\left(\lambda_i, N_i', \theta_i, N_i''\right)_{i\in\mathbb{N}}$ inductively as follows. We start by setting $\lambda_1 = 0$.

Suppose that we have already defined λ_i for some $i \in \mathbb{N}$. We use Proposition 3.5 to set $\theta_i = \theta_i\left(\varepsilon\right)$ and $N_i'' = N_i''\left(\varepsilon\right)$ such that

$$\mathbb{P}_N\left(\exists v \in B\left(KN\right) \text{ s.t. } diam\left(\mathcal{C}_a\left(v, \lambda_i\right)\right) \in \left[N, \left(1+\theta_i\right)N\right)\right) < \varepsilon 2^{-i-2}$$

for $N \geq N_i''$.

Suppose that we have already defined θ_i for some $i \in \mathbb{N}$. Then by Proposition 3.6 we set $\lambda_{i+1} = \lambda_{i+1}\left(\varepsilon\right)$ and $N_{i+1}' = N_{i+1}'\left(\varepsilon\right)$ such that the probability of the intersection of the events

- $\exists v \in B\left(KN\right)$ such that $diam\left(\mathcal{C}_a\left(v; \lambda_i\right)\right) \geq \left(1+\theta_i\right)N$, and
- $FC\left(p_{\lambda_i}\left(N\right), K+2, N\right) = FC\left(p_{\lambda_{i+1}}\left(N\right), K+2, N\right)$

is less than $2^{-i-2}\varepsilon$ for $N \geq N_{i+1}'$. Note that

$$\begin{aligned}\left\{\exists v \in B\left(KN\right) \text{ s.t. } diam\left(\mathcal{C}_a\left(v, \lambda_i\right)\right) \geq N\right\} &\supseteq \left\{\exists v \in B\left(\left(K+2\right)N\right) \text{ s.t. } diam\left(\mathcal{C}_a\left(v, \lambda_i\right)\right) \geq N\right\} \\ &= \left\{FC\left(p_{\lambda_i}\left(N\right), K+2, N\right) < FC\left(1, K+2, N\right)\right\}\end{aligned}$$

for $i \geq 1$. Hence the event

$$\left\{FC\left(p_{\lambda_i}(N), K+2, N\right) = FC\left(p_{\lambda_{i+1}}(N), K+2, N\right) < FC(1, K+2, N)\right\}$$

is a subset of the union of the events appearing in the definition of θ_i and λ_{i+1}. Thus the construction above gives that

$$\mathbb{P}_N\left(FC\left(p_{\lambda_i}(N), K+2, N\right) = FC\left(p_{\lambda_{i+1}}(N), K+2, N\right) < FC(1, K+2, N)\right) \leq 2^{-i-1}\varepsilon \tag{3.5}$$

for $i \geq 1$.

We set $N_0 = \bigvee_{i=1}^{L+1}\left(N_i' \vee N_i''\right)$. By (3.4) we have

$$\begin{aligned}
&\mathbb{P}_N(\text{ a cluster intersecting } B(KN) \text{ freezes after time } p_{\lambda_{L+1}}(N)) \\
&\quad= \mathbb{P}_N\left(FC\left(p_{\lambda_{L+1}}(N), K, N\right) < FC(1, K, N)\right) \\
&\quad\leq \mathbb{P}_N\left(FC\left(p_{\lambda_{L+1}}(N), K+2, N\right) \leq L,\ FC\left(p_{\lambda_{L+1}}(N), K+2, N\right) < FC(1, K+2, N)\right) + \mathbb{P}_N(L < F(1, K+2, N)) \\
&\quad\leq \varepsilon/2 + \mathbb{P}_N\left(\exists i \in [1, L+1] \cap \mathbb{Z} \text{ s.t. } FC\left(p_{\lambda_i}(N), K+2, N\right) = FC\left(p_{\lambda_{i+1}}(N, K+2, N)\right) < FC(1, K+2, N)\right) \\
&\quad\leq \varepsilon/2 + \sum_{i=1}^{L+1} \mathbb{P}_N\left(FC\left(p_{\lambda_i}(N, K+2, N)\right) = FC\left(p_{\lambda_{i+1}}(N, K+2, N)\right) < FC(1, K+2, N)\right) \\
&\quad\leq \varepsilon/2 + \sum_{i=1}^{L+1} 2^{-i-1}\varepsilon < \varepsilon
\end{aligned}$$

for $N \geq N_0$ where we applied (3.5) in the last line. This finishes the proof of Theorem 1.3. □

3.2 Proof of Corollary 1.6

Proof of Corollary 1.6 . For $\lambda \in \mathbb{R}$ and $N \in \mathbb{N}$ let $NF(\lambda) = NF(\lambda, N)$ denote the event that no cluster intersecting $B(5N)$ freezes after time $p_\lambda(N)$. By Theorem 1.3 there is $\lambda = \lambda(\varepsilon)$ and $N_1 = N_1(\varepsilon)$ such that

$$\mathbb{P}_N(NF(\lambda)) < \varepsilon/3 \tag{3.6}$$

for $N \geq N_1$.

First we consider the case where the origin is in an open frozen cluster at time 1, that is $diam(C(1)) \geq N$. Note that on the event $NF(\lambda)$, this frozen cluster was formed before or at $p_\lambda(N)$. Hence on this event there is a $p_\lambda(N)$-open path from the origin to distance at least $N/2$. Hence the event $\mathcal{A}_{1,0,o}^{\lambda,\lambda,N}(1, N/2)$ defined in Lemma 2.11 occurs.

Let us turn to the case where $diam(C(1)) < N$. Recall the notation $\mathcal{C}_a(\lambda)$ from Definition 3.4. It is easy to check $C(1) = \mathcal{C}_a(\lambda)$ that on the event$\{diam(C(1)) < N\} \cap NF(\lambda)$.

If $diam(\mathcal{C}_a(\lambda)) < aN$, then $\partial\mathcal{C}_a(\lambda) \cap B(2aN) \neq \emptyset$ for large N. Since $v \in \partial\mathcal{C}_a(\lambda) \cap B(2aN)$ is frozen, it has a neighbour which has an open frozen path to distance at least $N/2$. On the event $NF(\lambda)$, this path is $p_\lambda(N)$-open. Hence the event $\mathcal{A}_{1,0,o}^{\lambda,\lambda,N}(2aN, N/2)$ occurs. This combined with the argument above, for $a \in (0,1)$ and $N > N_2 = 1/a$ we have

$$\{diam(C(1)) \in [0, aN) \cup [N, \infty)\} \cap NF(\lambda) \subseteq \mathcal{A}_{1,0,o}^{\lambda,\lambda,N}(2aN, N/2).$$

Hence by Lemma 2.11 there is $c = c(\lambda)$ and $N_3 = N_3(\lambda)$ such that

$$\begin{aligned}
\mathbb{P}_N(diam(C(1)) \in [0, aN) \cup [N, \infty),\ NF(\lambda)) &\leq \mathbb{P}\left(\mathcal{A}_{1,0,o}^{\lambda,\lambda,N}(2aN, N/2)\right) \\
&\leq c\mathbb{P}_{1/2}(\mathcal{A}_{1,o}(2aN, N/2))
\end{aligned}$$

for $N \geq N_3$. Theorem 2.6 gives that there is $a = a(\varepsilon)$ and $N_4 = N_4(\varepsilon)$ such that

$$\mathbb{P}_N(diam(C(1)) \in [0, aN) \cup [N, \infty),\ NF(\lambda)) \leq c\mathbb{P}_{1/2}(\mathcal{A}_{1,o}(2aN, N/2)) < \varepsilon/3. \tag{3.7}$$

for $N \geq N_4$.

Finally, Proposition 3.5 gives $b = b(\varepsilon)$ and $N_5 = N_5(\varepsilon)$ such that

$$\begin{aligned}
\mathbb{P}_N(diam(\mathcal{C}_a(\lambda)) \in [bN, N),\ NF(\lambda)) &\leq \mathbb{P}_N(diam(\mathcal{C}_a(\lambda)) \in [bN, N)) \\
&\leq \varepsilon/3
\end{aligned} \tag{3.8}$$

for $N \geq N_5$.

Since $C(1) = \mathcal{C}_a(\lambda)$ on the event $\{diam(C(1)) < N\} \cap NF(\lambda)$, a combination of (3.6), (3.7) and (3.8) finishes the proof of Corollary 1.6. □

4 Proof of Proposition 3.6

4.1 Notation

Let us introduce some more notation. For $u = (u_1, u_2), v = (v_1, v_2) \in V$, we say that u is left (right) of v if $u_1 \leq v_1$ ($u_1 \geq v_1$). Similarly we say that u is below (above) v if $u_2 \leq v_2$ ($u_2 \geq v_2$). For a finite set of vertices $W \subseteq V$ we say that $v = (v_1, v_2) \in W$ is a leftmost (rightmost) vertex of W if for all $w = (w_1, w_2) \in W$, $v_1 \leq w_1$ ($v_1 \geq w_1$). We define the lowest and highest vertices of W in an analogous way.

Recall that $v, w \in V$, $v \sim w$ denotes that v and w are neighbours in $\mathbb{T}$. We extend this notation for subsets of V : For $S, U \subset V$, $S \sim U$ denotes that $\exists s \in S, \exists u \in U$ such that $s \sim u$.

Definition 4.1. Let $n \in \mathbb{N}$. We say that a sequence of vertices $v^1, v^2, \ldots, v^n$, denoted by ρ, is a path if

- $v^i \sim v^{i+1}$ for $i = 1, 2, \ldots, (n-1)$, and
- $v^i \neq v^j$ when $i \neq j$ for $i, j = 1, 2, \ldots, n$.

We say that ρ is non self touching, if $u, w \in \rho$ with $u \sim w$ then there is some $i \in \mathbb{N}$ with $1 \leq i \leq n-1$ such that either $u = v^i$ and $w = v^{i+1}$ or $u = v^{i+1}$ and $w = v^i$. We consider our paths to be ordered: v^1 is the starting point and v^n is the ending point of ρ. For $u, w \in \rho$ we say that u is after w in ρ, and denote it by $u \prec_\rho w$ if $u = v^i$ and $w = v^j$ for some $i, j \in \mathbb{N}$ with $1 \leq j \leq i \leq n$. For $u, w \in \rho$, $u \preceq_\rho w$ denotes that either $u = w$ or $u \prec_\rho w$. When it is clear from the context which path we are considering, we omit the subscript ρ. For $u, w, z \in \rho$ we say that w is in between u and z if $u \preceq w \preceq z$ or $u \succeq w \succeq z$. For $u, z \in \rho$ with $u \preceq z$ let $\rho_{u,z}$ denote the subpath of ρ consisting of the vertices between u and z.

We say that two paths ρ_1, ρ_2 are non-touching, if $\rho_1 \nsim \rho_2$.

Definition 4.2. Let $n \in \mathbb{N}$ and sequence of vertices $v^1, v^2, \ldots, v^n$, satisfying

- $v^i \sim v^{i+1 \mod n}$ for $i = 1, 2, \ldots, n$, and
- $v^i \neq v^j$ when $i \neq j$ for $i, j = 1, 2, \ldots, n$.

A loop ν is the equivalence class of the sequence $\left(v^1, v^2, \ldots, v^n\right)$ under cyclic permutations, i.e ν is the set of sequences $\left(v^j, v^{j+1 \mod n}, \ldots, v^{j+n-1 \mod n}\right)$ for $j = 1, 2, \ldots, n$. ν is non-self touching if for all $\left(w^1, w^2, \ldots, w^n\right) \in \nu$, the path $\left(w^1, w^2, \ldots, w^{n-1}\right)$ is non-self touching.

We say that ν contains a vertex v and denote it by $v \in \nu$ if $v = v^i$ for some $i \in \{1, 2, \ldots, n\}$. Let $v, w \in \nu$ with $v \neq w$ and let ρ denote the unique path which starts at v and represents ν. With the notation of Definition 4.1, let $\nu_{v,w} := \rho_{v,w}$ denote the arc of ν starting at v and ending at w.

4.2 Thick paths

Definition 4.3. Let $M \in \mathbb{N}$ be fixed. The M-grid is the set of parallelograms $B\left((2M+1)z; M\right)$ for $z \in V$. Let π be a sequence consisting of some parallelograms of the M-grid. We say that π is an M-gridpath, if for any two consecutive parallelograms B, B' of π share a side, i.e $|\partial B \cap B'| \geq 2$.

Definition 4.4. Let C be a subgraph of $\mathbb{T}$, $D \subset V$ and $a, b \in \mathbb{N}$. We say that C is (a,b)-nice in D, if it satisfies the conditions

1. C is a connected induced subgraph of $\mathbb{T}$,
2. ∂C is a disjoint union of non-touching loops, each with diameter bigger than $2b$.
3. Let $u, v \in \partial C \cap D$ with $d(u,v) \leq a$. Then u, v are contained in the same loop γ of ∂C, and $diam\left(\gamma_{u,v}\right) \wedge diam\left(\gamma_{v,u}\right) \leq b$.

In the case where $B = V$, we say that C is (a,b)-nice.

Let C be (a,b)-nice for some $a, b \in \mathbb{N}$. Condition 3 of Definition 4.4, roughly speaking, says that if there is a corridor in C with width less than a, then it connects two parts of C such that one part has diameter at most b. This suggests that when b is small compared to $diam(C)$, when we can move a parallelogram with side length $O(a)$ in C between two distant points of C. This intuitive argument leads us to the following lemma.

Lemma 4.5. *Let $a, b \in \mathbb{N}$ with $a \geq 2000$. Let C be an (a,b)-nice subgraph of $\mathbb{T}$. Then there is a $\lfloor a/200 - 10 \rfloor$-gridpath contained in C with diameter at least $diam\,(C) - 2b - 2a - 12$.*

We use the following 'local' version of Lemma 4.5:

Lemma 4.6. *Let $a, b, c \in \mathbb{N}$ with $a \geq 2000$. Let C be subgraph of $\mathbb{T}$ which is (a,b)-nice in $B\,(c)$. Let C' be a connected component of $C \cap B\,(c)$. Then there is a $\lfloor a/200 - 10 \rfloor$-gridpath contained in C' with diameter at least $diam\,(C') - 2b - 2a - 12$.*

Proof of Lemma 4.5 and 4.6. The proof of Lemma 4.5 and 4.6 have geometric/topologic nature, hence it is moved to Section A.2 of the Appendix. ☐

We recall and prove Proposition 3.6 in the following.

Proposition 3.6. *Let $\theta \in (0,1)$, $\varepsilon > 0$ and $\lambda_1 K, \in \mathbb{R}$. Recall the notation $FC\,(t, K+2, N)$ from Lemma 3.3. There exists $\lambda_2 = \lambda_2\,(\lambda_1, \theta, \varepsilon)$ and $N_0 = N_0\,(\lambda_1, \theta, \varepsilon)$ such that the probability of the intersection of the events*

- *$\exists v \in B\,(KN)$ such that $diam\,(\mathcal{C}_a\,(v; \lambda_1, N)) \geq (1+\theta)\,N$, and*
- *none of the clusters intersecting $B\,((K+2)\,N)$ freeze in the time interval $(p_{\lambda_1}\,(N), p_{\lambda_2}\,(N)]$, i.e.*

$$FC\,(p_{\lambda_1}\,(N), K+2, N) = FC\,(p_{\lambda_2}\,(N), K+2, N)$$

is less than ε for $N \geq N_0$.

Proof of Proposition 3.6. By Lemma 2.27 we choose $\lambda_0 = \lambda_0\,(\varepsilon, K) \leq \lambda_1$ and $N_1 = N_1\,(\varepsilon, K)$ such that

$$\mathbb{P}\,(\mathcal{N}_c\,(\lambda_0, 1/6, K+6, N)) > 1 - \varepsilon/3. \tag{4.1}$$

By Corollary 2.13 we choose $\eta < \theta/10$ and $N_2 = N_2\,(\eta, \theta, \lambda_0, \lambda_1, K)$ such that

$$\mathbb{P}\,(\mathcal{NA}\,(2\eta, \theta/10, \lambda_0, \lambda_1, K+4, N)) > 1 - \varepsilon/3 \tag{4.2}$$

for all $N \geq N_2$. Let

$$E := \mathcal{N}_c\,(\lambda_0, 1/6, K+6, N) \cap \mathcal{NA}\,(2\eta, \theta, \lambda_0, \lambda_1, K+4, N)\,.$$

Claim 4.7. Let $u \in B\,(KN)$ with $diam\,(\mathcal{C}_a\,(u; \lambda_1, N)) \geq (1+\theta)\,N$. Then $\mathcal{C}_a\,(u; \lambda_1, N)$ is $\left(\eta N, \frac{\theta}{10} N\right)$-nice in $B\,(u; 2N)$ on the event E.

Proof. Let us check the conditions of Definition 4.4. The Condition 1 is satisfied by the definition of $\mathcal{C}_a\,(u; \lambda_1, N)$.

All the holes of $\mathcal{C}_a\,(u; \lambda_1, N)$ contain a frozen cluster, which have diameter at least N. This combined with $2\frac{\theta}{10}N < N$, shows that Condition 2 of Definition 4.4 holds.

Let $x, y \in \partial\mathcal{C}_a\,(u; \lambda_1, K) \cap B\,(u; 2N)$ with $d\,(x,y) \leq \eta N$. We have two cases.

Case 1. x, y lie in different loops of $\partial\mathcal{C}_a\,(u; \lambda_1, N)$. For $i = x, y$, let γ_i denote the loop containing i. Furthermore, let $\tilde{\gamma}_i$ denote the connected component of i in $\gamma_i \cap B\,(i; 2N)$. We have $diam\,(\tilde{\gamma}_i) \geq N$. Moreover, $\tilde{\gamma}_i \subset B\,(i; 2N) \subset B\,((K+4)\,N)$. Observation 3.2 gives that on the event $\mathcal{N}_c\,(\lambda_0, 1/6, K+6, N)$, $\tilde{\gamma}_i$ is $p_{\lambda_0}\,(N)$-closed. Hence each of $\tilde{\gamma}_x$ and $\tilde{\gamma}_y$ gives two closed $p_{\lambda_0}\,(N)$-closed arms in $A\,(x; 2\eta N, N/2)$. Moreover, the frozen clusters neighbouring x and y provide two disjoint $p_{\lambda_1}\,(N)$-open arms. Hence there are 6 disjoint arms in $A\,(x; 2\eta N, N/2)$, thus $\mathcal{NA}^c\,(2\eta, \theta/10, \lambda_0, \lambda_1, K+4, N)$ occurs.

Case 2. x, y lie on the same loop of $\partial\mathcal{C}_a\,(u; \lambda_1, N)$. This case can be treated similarly Condition 1, with the difference that if x, y violate Condition 3 of Definition 4.4 then we get 6 arms in $A\left(x; 2\eta N, \frac{\theta}{10}N\right)$. Hence $\mathcal{NA}^c\,(2\eta, \theta/10, \lambda_0, \lambda_1, K+4, N)$ occurs.

Hence in both cases E^c occurs. Thus on the event E all the conditions of Definition 4.4 are satisfied for $\mathcal{C}_a\,(u; \lambda_1, N)$, which finishes the proof of Claim 4.7. ☐

Let us turn back to the proof of Proposition 3.6. Let $u \in B\,(KN)$ with $diam\,(\mathcal{C}_a\,(u; \lambda_1, N)) \geq (1+\theta)\,N$. Let $\tilde{\mathcal{C}}_a\,(u, \lambda_1, N)$ denote the connected component of u in $\mathcal{C}_a\,(u, \lambda_1, N) \cap B\,(2N)$. Since $diam\,(\mathcal{C}_a\,(u; \lambda_1, N)) \geq (1+\theta)\,N$ and $\theta < 1$, we have $diam\left(\tilde{\mathcal{C}}_a\,(u; \lambda_1, N)\right) \geq (1+\theta)\,N$. By Lemma 4.6 we set $\eta = \eta\,(\theta) \in (0, \theta/100)$ and $N_3 = N_3\,(\theta)$ such that on the event E for all $u \in B\,(KN)$, with $diam\,(\mathcal{C}_a\,(u; \lambda_1, N)) \geq (1+\theta)\,N$ there is a $\lfloor \eta N \rfloor$-gridpath $\rho_u \subset \tilde{\mathcal{C}}_a\,(u; \lambda_1, N)$ with $diam\,(\rho_u) \geq (1+\theta/2)\,N$ for $N \geq N_3$.

Lemma 2.27 gives that there is $\lambda_2 = \lambda_2(\varepsilon, \eta, K)$ and $N_4 = N_4(\varepsilon, \eta, K)$ such that

$$\mathbb{P}\left(\mathcal{N}_o\left(\lambda_2, \eta/2, K+4, N\right)\right) > 1 - \varepsilon/3 \tag{4.3}$$

for $N \geq N_4(\varepsilon, \eta, K)$. We set $N_0 := \bigvee_{i=1}^{4} N_i$. Let

$$\begin{aligned} G :=& E \cap \mathcal{N}_o\left(\lambda_2, \eta/2, K+4, N\right), \\ M :=& \left\{\exists v \in B(KN) \text{ s.t. } \mathit{diam}\left(\mathcal{C}_a\left(v; \lambda_1, N\right)\right) \geq (1+\theta) N\right\} \cap G. \end{aligned}$$

Combination of (4.1), (4.2) and (4.3) gives that

$$\mathbb{P}\left(G^c\right) < \varepsilon \tag{4.4}$$

for $N \geq N_0$.

Recall that for $N \geq N_0$, on the event E for $u \in B(KN)$, with $\mathit{diam}\left(\mathcal{C}_a\left(u; \lambda_1, N\right)\right) \geq (1+\theta) N$ there is a $\lfloor \eta N \rfloor$-gridpath $\rho_u \subset \tilde{\mathcal{C}}_a\left(u; \lambda_1, N\right)$ with $\mathit{diam}\left(\rho_u\right) \geq (1+\theta/2) N$. On the event $\mathcal{N}_o\left(\lambda_2, \eta/2, K+4, N\right)$, this gridpath $\rho_u \subseteq B((K+2)N)$ contains a $p_{\lambda_2}(N)$-open component with diameter at least N. Hence on the event M, at least one cluster intersecting $B((K+2)N)$ freezes in the time interval $\left(p_{\lambda_1}(N), p_{\lambda_2}(N)\right]$.That is

$$M \subseteq \left\{FC\left(p_{\lambda_1}(N), K+2, N\right) < FC\left(p_{\lambda_2}(N), K+2, N\right)\right\}.$$

Thus

$$\left\{\exists v \in B(KN) \text{ s.t. } \mathit{diam}\left(\mathcal{C}_a\left(v; \lambda_1, N\right)\right) \geq (1+\theta) N\right\} \cap \left\{FC\left(p_{\lambda_1}(N), K+2, N\right) = FC\left(p_{\lambda_2}(N), K+2, N\right)\right\} \subset G^c,$$

which together with (4.4) finishes the proof of Proposition 3.6. □

5 Proof of Proposition 3.5

5.1 Lowest point of the lowest crossing in parallelograms

The central object of this section is the following. Recall the notation of Section 4.1.

Definition 5.1. Let R be a connected subgraph of $\mathbb{T}$ and let $r \subset \partial R$. We define $\mathcal{L}(R, r)$ as the the (random) set of lowest vertices $v \in R$ such that v is closed, and there are two non-touching closed paths in R starting at a vertex neighbouring to v and ending at r.

Consider the site percolation model on the triangular lattice with parameter $p \in [0,1]$. We investigate the distribution of $\mathcal{L}(R, r)$ in the case where $p = p_\lambda(N)$, $R = B(bN)$ and $r = \mathit{top}(B(bN)) := [-bN, bN] \boxtimes \{\lfloor bN \rfloor + 1\}$ for $\lambda \in \mathbb{R}$ and $b > 0$.

The distribution of $\mathcal{L}(R, r)$ was examined in [22] in the case where $p = 1/2$, R is the upper half plane and $r = [0, n] \boxtimes \{0\}$ for some $n \in \mathbb{N}$. The results of [22] do not imply, nor follow from ours.

Definition 5.2. For a parallelogram B, let $HCr(B)$ denote set of paths in B which connect the left and the right sides of B. For $\rho \in HCr(B)$, let $Be(\rho) = Be(\rho, B)$ denote the set of vertices in B which are 'under' ρ. It is the set of vertices $v \in B \setminus \rho$ which are connected to the bottom side of $B(bN)$. Furthermore, we define $Ab(\rho) = Ab(\rho, B) := B \setminus (\rho \cup Be(\rho, B))$.

Lemma 5.3. *Let $a, b \in (0,1)$ with $5a < b$ and $\lambda \in \mathbb{R}$. For $k, l, N \in \mathbb{N}$ with $l < k$ we define the event*

$$L_{l,k} = \left\{\mathcal{L}\left(B(bN), \mathit{top}(bN)\right) \cap B_{l,k} \neq \emptyset\right\}.$$

In words this is the event that at least one of the lowest vertices of $B(bN)$ with two non-touching closed paths $B(bN)$ to the top side of $B(bN)$ is in the parallelogram

$$B_{l,k} = [-aN, aN] \boxtimes \left(\left(2\frac{l}{k} - 1\right) aN, \left(2\frac{l+1}{k} - 1\right) aN\right].$$

Let $\lambda_1, \lambda_2 \in \mathbb{R}$. Then there exist $C = C(a, b, \lambda_1, \lambda_2)$ and $N_0 = N_0(a, b, \lambda_1, \lambda_2, k)$ such that for all $\lambda \in [\lambda_1, \lambda_2]$ and $k, l \in \mathbb{N}$ with $l \leq k-1$ we have

$$\mathbb{P}_{p_\lambda(N)}\left(L_{l,k}\right) \leq C k^{-1} \tag{5.1}$$

for $N \geq N_0$.

Proof of Lemma 5.3. Note that $L_{l,k} \cap L_{m,k} = \emptyset$ when $l \neq m$. Hence it is enough to show that there is exists $c = c(a, b, \lambda_1, \lambda_2) > 0$ and $N_0 = N_0(a, b, \lambda_1, \lambda_2)$ such that for all $l, m \in [0, k-1] \cap \mathbb{Z}$ we have

$$c\mathbb{P}_{p_\lambda(N)}(L_{l,k}) \leq \mathbb{P}_{p_\lambda(N)}(L_{m,k}) \tag{5.2}$$

for $N \geq N_0$.

Let k, l be given. Let s_L (s_R) denote the left (right) endpoint of $top(bN)$. We say that a path $\rho \subseteq B(bN) \cup top(bN)$ is good, if it

- starts at s_L and ends at s_R,
- it is non-self touching
- and one of its lowest points is in $B_{l,k}$.

Let ρ be some given good path. Recall Definition 5.2 and let $Be(\rho) = Be(\rho, (B(bN)))$. Let H_ρ denote the event that there are two open paths in $Be(\rho) \cap [-bN, bN] \boxtimes \left[aN, \frac{b-2a}{2}N\right]$ from the left and right sides of the parallelogram $[-bN, bN] \boxtimes \left[aN, \frac{b-2a}{2}N\right]$ to ρ. Let γ denote the lowest non-self touching path in $B(bN) \cup top(bN)$ which starts at s_L and ends at s_R, and of which all the vertices outside of $top(bN)$ are closed. On the event $L_{l,k}$ γ is good.

Case 1. First we consider the case where $l < m$. Recall the definition of increasing events from Definition 2.2. For any good path ρ, on the event that all of the vertices of $\rho \setminus top(bN)$ are closed, the event $\{\gamma = \rho\}$ is increasing in the configuration in $B(bN) \setminus \rho$. Let $H_1 := \mathcal{H}_o([-bN, bN] \boxtimes [-bN, -(b-a)N])$. The events H_1 and H_ρ are increasing. Thus by FKG and by Lemma 2.20, we have

$$\begin{aligned}
&\mathbb{P}_{p_\lambda(N)}(L_{l,k} \cap H_1 \cap H_\gamma) \\
&\quad = \sum_{\rho \text{ good}} \mathbb{P}_{p_\lambda(N)}(H_1 \cap H_\rho \cap \{\gamma = \rho\} \mid \rho \setminus top(bN) \text{ is closed })\,\mathbb{P}_{p_\lambda(N)}(\rho \setminus top(bN) \text{ is closed}) \\
&\quad \geq \sum_{\rho \text{ good}} \mathbb{P}_{p_\lambda(N)}(\{\gamma = \rho\} \mid \rho \setminus top(bN) \text{ is closed})\,\mathbb{P}_{p_\lambda(N)}(H_1 \mid \rho \setminus top(bN) \text{ is closed}) \\
&\qquad\qquad \mathbb{P}_{p_\lambda(N)}(H_\rho \mid \rho \setminus top(bN) \text{ is closed})\,\mathbb{P}_{p_\lambda(N)}(\rho \setminus top(bN) \text{ is closed}) \\
&\quad \geq \sum_{\rho \text{ good}} c_1\mathbb{P}_{p_\lambda(N)}(\{\gamma = \rho\} \mid \rho \setminus top(bN) \text{ is closed})\,\mathbb{P}_{p_\lambda(N)}(\rho \setminus top(bN) \text{ is closed}) \\
&\quad = c_1\mathbb{P}_{p_\lambda(N)}(L_{l,k})
\end{aligned} \tag{5.3}$$

for $c_1 = c_1(a, b, \lambda_1, \lambda_2) > 0$ and for $N \geq N_1 = N_1(a, b, \lambda_1, \lambda_2)$.

There is $N_2 = N_2(k)$ such that for $N \geq N_2$ and for all $l, m \in [0, k-1] \cap \mathbb{Z}$ with $l < m$ there is a shift $S = S(l, m, k)$ which moves the parallelogram $B_{l,k}$ to a subset of $B_{m-1,k} \cup B_{m,k}$. Let us take a configuration in $\omega \in \{o, c\}^V$ for which the event $L_{l,k} \cap H_1 \cap H_\gamma$ holds. Then shifted configuration $S(\omega)$ satisfies $L_{m-1,k} \cup L_{m,k}$. Hence

$$\begin{aligned}
\mathbb{P}_{p_\lambda(N)}(L_{m-1,k} \cup L_{m,k}) &\geq \mathbb{P}_{p_\lambda(N)}(L_{l,k} \cap H_1 \cap H_\gamma), \\
&\geq c_1\mathbb{P}_{p_\lambda(N)}(L_{l,k})
\end{aligned} \tag{5.4}$$

by (5.3) for $N \geq N_1 \vee N_2$.

Case 2. When $l > m$ we have a similar argument. The difference is that now we want to shift downwards. To get a configuration in $L_{m,k}$ after the shift, we have to extend the closed path γ see Figure 1 for more details.

Let ρ be a fixed good path. Let O_ρ denote the event that there is path ν such that

- $\nu \subseteq B_0 := [-bN, bN] \boxtimes \left[-bN, \frac{b}{4}N\right]$,
- ν connects the left and the right sides of the parallelogram $B_1 := [-bN, bN] \boxtimes \left[aN, \frac{b}{4}N\right]$,
- ν is a concatenation of some open paths which lie in $Be(\rho) \cap B_1$, and of some subpaths of ρ.

Clearly, O_ρ is an increasing event. On O_ρ, let $\xi(\rho)$ denote the lowest path which satisfies the conditions in the definition of O_ρ. Recall the definition of decreasing events from Definition 2.2, and the definition of γ from Case 1. Let us condition on the event that all the vertices of $\rho \setminus top(bN)$ are closed. Then the event $\{\gamma = \rho\}$ is increasing on the configuration in $B(bN)\setminus\rho$, and it only depends on the configuration in $Be(\rho)$. Hence a combination of FKG and Corollary 2.20 give that

$$\begin{aligned}
&\mathbb{P}_{p_\lambda(N)}\left(L_{l,k}\cap O_\gamma\right)\\
&\quad=\sum_{\rho\text{ good}}\mathbb{P}_{p_\lambda(N)}\left(O_\rho\cap\{\gamma=\rho\}\mid\rho\setminus top(bN)\text{ is closed}\right)\mathbb{P}_{p_\lambda(N)}\left(\rho\setminus top(bN)\text{ is closed}\right)\\
&\quad\geq\sum_{\rho\text{ good}}\mathbb{P}_{p_\lambda(N)}\left(\{\gamma=\rho\}\mid\rho\setminus top(bN)\text{ is closed}\right)\mathbb{P}_{p_\lambda(N)}\left(O_\rho\mid\rho\setminus top(bN)\text{ is closed}\right)\mathbb{P}_{p_\lambda(N)}\left(\rho\setminus top(bN)\text{ is closed}\right)\\
&\quad\geq\sum_{\rho\text{ good}}\mathbb{P}_{p_\lambda(N)}\left(\{\gamma=\rho\}\mid\rho\setminus top(bN)\text{ is closed}\right)\mathbb{P}_{p_\lambda(N)}\left(\mathcal{H}_o(B_1)\right)\mathbb{P}_{p_\lambda(N)}\left(\rho\setminus top(bN)\text{ is closed}\right)\\
&\quad\geq c_2(\lambda_1,\lambda_2,a,b)\,\mathbb{P}_{p_\lambda(N)}\left(L_{l,k}\right)
\end{aligned}\tag{5.5}$$

for $c_2=c_2(a,b,\lambda_1,\lambda_2)>0$ and for $N\geq N_3=N_3(a,b,\lambda_1,\lambda_2)$.

For $W\subseteq V$ and $\omega\in\{o,c\}^V$, $\omega_W\in\{o,c\}^W$ denotes the restriction of ω to the configuration in W. That is $\omega_W(v)=\omega(v)$ for $v\in W$. Recall Definition 5.2. Let $\zeta\in HCr(B_0)$ be arbitrary. It is easy to check that the event $L_{l,k}\cap O_\gamma\cap\{\xi(\gamma)=\zeta\}$ is decreasing in the configuration in $Ab(\zeta)$. Let us take the parallelograms $B_2=[-bN,bN]\boxtimes\left[\frac{b}{4}N,\frac{b}{2}N\right]$, $B_3=[-bN,bN]\boxtimes\left[\frac{3}{4}bN,bN\right]$, $B_4=\left[-bN,-\frac{1}{2}bN\right]\boxtimes\left[\frac{1}{4}bN,(b+2a)N\right]$ and $B_5=\left[\frac{1}{2}bN,bN\right]\boxtimes\left[\frac{1}{4}bN,(b+2a)N\right]$. Let $\mathcal{D}=\mathcal{H}_c(B_2)\cap\mathcal{H}_c(B_3)\cap\mathcal{V}_c(B_4)\cap\mathcal{V}_c(B_5)$. Clearly, $\mathcal{D}$ is a decreasing event. Hence a combination of FKG and Corollary 2.20 give that for $c_3=c_3(a,b,\lambda_1,\lambda_2)>0$ and $N\geq N_4=N_4(a,b,\lambda_1,\lambda_2)$ we have

$$\begin{aligned}
&\mathbb{P}_{p_\lambda(N)}\left(L_{k,l}\cap O_\gamma\cap\mathcal{D}\right)\\
&\quad=\sum_\zeta\sum_\sigma\mathbb{P}_{p_\lambda(N)}\left(L_{k,l}\cap O_\gamma\cap\{\xi(\gamma)=\zeta\}\cap\mathcal{D}\mid\omega_{\zeta\cup Be(\zeta)}=\sigma\right)\mathbb{P}_{p_\lambda(N)}\left(\omega_{\zeta\cup Be(\zeta)}=\sigma\right)\\
&\quad\geq\sum_\zeta\sum_\sigma\mathbb{P}_{p_\lambda(N)}\left(L_{k,l}\cap O_\gamma\cap\{\xi(\gamma)=\zeta\}\mid\omega_{\zeta\cup Be(\zeta)}=\sigma\right)\mathbb{P}_{p_\lambda(N)}\left(\mathcal{D}\mid\omega_{\zeta\cup Be(\zeta)}=\sigma\right)\mathbb{P}_{p_\lambda(N)}\left(\omega_{\zeta\cup Be(\zeta)}=\sigma\right)\\
&\quad=\sum_\zeta\sum_\sigma\mathbb{P}_{p_\lambda(N)}\left(L_{k,l}\cap O_\gamma\cap\{\xi(\gamma)=\zeta\}\mid\omega_{\zeta\cup Be(\zeta)}=\sigma\right)\mathbb{P}_{p_\lambda(N)}(\mathcal{D})\,\mathbb{P}_{p_\lambda(N)}\left(\omega_{\zeta\cup Be(\zeta)}=\sigma\right)\\
&\quad\geq c_3(a,b,\lambda_1,\lambda_2)\sum_\zeta\sum_\sigma\mathbb{P}_{p_\lambda(N)}\left(L_{k,l}\cap O_\gamma\cap\{\xi(\gamma)=\zeta\}\mid\omega_{\zeta\cup Be(\zeta)}=\sigma\right)\mathbb{P}_{p_\lambda(N)}\left(\omega_{\zeta\cup Be(\zeta)}=\sigma\right)\\
&\quad=c_3(a,b,\lambda_1,\lambda_2)\,\mathbb{P}_{p_\lambda(N)}\left(L_{k,l}\cap O_\gamma\right)
\end{aligned}\tag{5.6}$$

where the summation in ζ is over $HCr(B_0)$ and the summation in σ is over $\{o,c\}^{\zeta\cup Be(\zeta)}$. In the third line we used that $\mathcal{D}$ does not depend on the configuration in $\zeta\cup Be(\zeta)$.

There is $N_5=N_5(k)$ such that for $N\geq N_5$ and for all $l,m\in[0,k-1]\cap\mathbb{Z}$ with $l>m$ there is a shift $S=S(l,m,k)$ which moves the parallelogram $B_{l,k}$ to a subset of $B_{m,k}\cup B_{m+1,k}$. Let us take a configuration $\omega\in\{o,c\}^V$ which satisfies $L_{k,l}\cap O_\gamma\cap\mathcal{D}$. Then the shifted configuration $S(\omega)$ satisfies $L_{m,k}\cup L_{m+1,k}$, see Figure 1 for more details. Hence for $N\geq N_3\vee N_4\vee N_5$ we have

$$\begin{aligned}
\mathbb{P}_{p_\lambda(N)}\left(L_{m,k}\cup L_{m+1,k}\right)&\geq\mathbb{P}_{p_\lambda(N)}\left(L_{k,l}\cap O_\gamma\cap\mathcal{D}\right)\\
&\geq c_2c_3\mathbb{P}_{p_\lambda(N)}\left(L_{l,k}\right)
\end{aligned}\tag{5.7}$$

by a combination of (5.5) and (5.6).

(5.4) together with (5.7) implies (5.2) and finishes the proof of Lemma 5.3. □

Remark 5.4. The proof of the lemma shows that $\mathbb{P}_{p_\lambda(N)}\left(L_{l,k}\right)=O\left(k^{-1}\right)$: Let $a,b,\lambda,\lambda_1,\lambda_2$ be as in Lemma 5.3. Standard RSW techniques give that there is $c'=c'(a,b,\lambda_1,\lambda_2)>0$ and $N_0=N_0(a,b,\lambda_1,\lambda_2)$ such that

$$\mathbb{P}_{p_\lambda(N)}\left(\mathcal{L}(B(bN),t(bN))\cap B(aN)\neq\emptyset\right)\geq c'$$

for $N\geq N_0$. This combined with the proof of Lemma 5.3, we get that some $C'=C'(a,b,\lambda_1,\lambda_2)>0$ and $N_1=N_1(a,b,\lambda_1,\lambda_2,k)$ we have

$$\mathbb{P}_{p_\lambda(N)}\left(L_{l,k}\right)\geq C'k^{-1}$$

for $N\geq N_1$.



Figure 1: The continuous line represents γ. The dashed paths are the closed crossings of $\mathcal{D}$, which allow us to prolong γ. The dashed-dotted paths are the open parts of $\xi(\gamma)$. They, together with γ, prevent the occurrence of closed vertices below the lowest point of γ with two closed arms to the top side after the shift.

5.2 Lowest point of the lowest crossing in regular regions

Recall Definition 5.1. Let $B \subset B'$ be parallelograms, and let R be a subgraph of $\mathbb{T}$ with $B \subset R \subset B'$. Furthermore let $r \subset \partial R$. Our next aim is to compare the event $\mathcal{L}(R, r) \cap B \neq \emptyset$ to $\mathcal{L}(B', top(B')) \cap B \neq \emptyset$ in the case where the pair (R, r) is 'regular'. We make this precise in the following.

We say that an subgraph $H \subseteq \mathbb{T}$ is simply connected, if it is connected and for all loops $\sigma \subseteq H$, all of the finite components of $\mathbb{T} \setminus \sigma$ are contained in H.

Definition 5.5. Let $a, b \in \mathbb{N}$ such that $5a < b$. A pair (R, r) is (a, b)-regular, if

1. R is an induced subgraph of $\mathbb{T}$ such that $cl(R)$ is simply connected,
2. $B(a) \subseteq R \subseteq B(b)$,
3. $r \subset \partial R$, such that $\emptyset \neq r \subsetneqq \partial R$. Furthermore, r and $\partial R \setminus r$ are self-avoiding paths such that R is on the right hand side, as we walk along them.
4. $r \subseteq [-b, b] \boxtimes [5a, b]$.

Lemma 5.6. *Let $a, b \in (0, 1)$ with $5a < b$ and $\lambda \in \mathbb{R}$. Let (R, r) be an (aN, bN)-regular graph. For $k, l, N \in \mathbb{N}$ with $l < k$ we define the events*

$$L_{l,k}(B(2bN), top(B(2bN))) := \{\mathcal{L}(B(2bN), top(2bN)) \cap B_{l,k} \neq \emptyset\},$$
$$L_{l,k}(R, r) := \{\mathcal{L}(R, r) \cap B_{l,k} \neq \emptyset\},$$

where

$$B_{l,k} := [-aN, aN] \boxtimes \left(\left(2\frac{l}{k} - 1\right) aN, \left(2\frac{l+1}{k} - 1\right) aN\right].$$

Let $\lambda_1, \lambda_2 \in \mathbb{R}$. Then there exist $C = C(a, b, \lambda_1, \lambda_2)$ and $N_0 = N_0(a, b, \lambda_1, \lambda_2, k)$ such that for all $\lambda \in [\lambda_1, \lambda_2]$ and $k, l \in \mathbb{N}$ with $l \leq k - 1$ we have

$$\mathbb{P}_{p_\lambda(N)}(L_{l,k}(R, r)) \leq C\mathbb{P}_{p_\lambda(N)}(L_{l,k}(B(2bN), top(B(2bN)))) \tag{5.8}$$

for $N \geq N_0$.



Figure 2: The dashed paths are the closed crossings of the event $\mathcal{D}$ which allow us to prolong γ. The dashed-dotted paths are the open parts of $\xi(\gamma)$. They, together with γ, prevent the occurrence of closed vertices below the lowest point of γ with two closed arms to the top side of $B(2bN)$.

Proof of Lemma 5.6. The proof follows the arguments of Case 2 in the proof of Lemma 5.3. Our aim is to show that, conditioning on $L_{l,k}(R,r)$, the open and closed crossings of Figure 2 occur with probability bounded away from 0 cf. Figure 1.

Let k,l be given. Let s_L (s_R) denote the starting (ending) vertex of r. We say that a path $\rho \subseteq R \cup r$ is good, if it

- starts at s_L and ends at s_R,
- it is non-self touching
- and one of its lowest points is in $B_{l,k}$.

Let ρ be a fixed good path. Let $Be(\rho, R)$ denote the set of vertices in R ‘under‘ ρ. It is the intersection of R with the connected component of $\partial R \setminus r$ in $cl(R) \setminus r$. Let $Ab(\rho, R) := R \setminus Be(\rho, R)$. Recall Definition 5.2.

Let O_ρ denote the event that there is path ν such that

- ν is non self-touching,
- $\nu \subseteq B_0 := [-2bN, 2bN] \boxtimes [-aN, 2aN]$,
- ν connects the left and the right side of the parallelogram $B_1 := [-2bN, 2bN] \boxtimes [aN, 2aN]$,
- $v \setminus R \subset B_1$ and the vertices in $\nu \setminus R$ are open,
- each of the paths of $\nu \cap R$ is a concatenation of some open paths which lie in $Be(\rho, B(bN)) \cap B_1$, and of some subpaths of ρ.

Let γ denote the lowest non-self touching path in $R \cup r$ which starts at s_L and ends at s_R, and of which all the vertices outside of r are closed. Note that on the event $L_{l,k}(R,r)$, γ is good. By simple modifications of the arguments of Case 2 in the proof of Lemma 5.3 we get that there are $c_1 = c_1(a,b,\lambda_1,\lambda_2) > 0$ and $N_1 = N_1(a,b,\lambda_1,\lambda_2)$ such that

$$\mathbb{P}_{p_\lambda(N)}\left(L_{l,k}(R,r) \cap O_\gamma\right) \geq c_1 \mathbb{P}_{p_\lambda(N)}\left(L_{l,k}(R,r)\right) \tag{5.9}$$

for $l,k \in \mathbb{N}$, $0 \leq l \leq k-1$, $\lambda \in [\lambda_1, \lambda_2]$ for $N \geq N_1$.

Recall Definition 5.2. Let $\zeta \in HCr(B(2bN))$. On the event $L_{l,k}(R,r) \cap O_\gamma$ we have $R \cap (\mathbb{Z} \boxtimes [3aN, bN]) \subset Ab(\xi(\gamma), B(2bN))$. Hence the event $L_{l,k}(R,r) \cap O_\gamma \cap \{\xi(\gamma) = \zeta\}$ is decreasing on the configuration in $Ab(\zeta, B(2bN))$.

Let $B_2 = [-2bN, 2bN] \boxtimes [3aN, 4aN]$, $B_3 = [-2bN, -bN] \boxtimes [3aN, 2bN]$, $B_4 = [bN, 2bN] \boxtimes [3aN, 2bN]$ and $\mathcal{D} = \mathcal{H}_c(B_2) \cap \mathcal{V}_c(B_3) \cap \mathcal{V}_c(B_4)$. The arguments of Case 2 of Lemma 5.3 give that there exist $c_2 = c_2(a, b, \lambda_1, \lambda_2) > 0$ and $N_2 = N_2(a, b, \lambda_1, \lambda_2, k)$ such that

$$\mathbb{P}_{p_\lambda(N)}(L_{l,k}(R, r) \cap O_\gamma \cap \mathcal{D}) \geq c_2 \mathbb{P}_{p_\lambda(N)}(L_{l,k}(R, r) \cap O_\gamma) \tag{5.10}$$

for $l, k \in \mathbb{N}$, $0 \leq l \leq k-1$, $\lambda \in [\lambda_1, \lambda_2]$ for $N \geq N_2$. Note that $L_{l,k}(R, r) \cap O_\gamma \cap \mathcal{D} \subset L_{l,k}(B(2bN), top(2bN))$. See Figure 2 for more details. This combined with (5.9) and (5.10) finishes the proof of Lemma 5.6. □

Remark 5.7. We could prove a version of Lemma 5.6 where we compare the probability of the events $L_{l,k}(R, r)$ and $L_{l,k}(B(bN, top(B(bN))))$. In view of Lemma 5.3, these two versions are equivalent. However, in the proof of this alternative version, similarly to Case 2 in the proof of Lemma 5.3, we would need an extra horizontal crossing to prolong the closed arms of γ. This would complicate Figure 2. This is the reason why we chose to prove Lemma 5.6 in its current form.

A combination of Lemma 5.3 and 5.6 gives the following:

Corollary 5.8. *Suppose that the conditions of Lemma 5.3 hold. Then there exist* $c = c(a, b, \lambda_1, \lambda_2)$ *and* $N_0 = N_0(a, b, \lambda_1, \lambda_2, k)$ *such that*

$$\mathbb{P}_{p_\lambda(N)}(L_{l,k}(R, r)) \leq ck^{-1}$$

for $l = 0, 1, \ldots, k-1$, $\lambda \in [\lambda_1, \lambda_2]$ *and* $N \geq N_0$.

5.3 The diameter of the active clusters close to time 1/2

We turn to the N-parameter frozen percolation process. In the introduction we indicated that the N-parameter frozen percolation process exists since it is a finite range interacting particle system. It is also true that the process is measurable with respect to the τ values. The following lemma follows from the arguments in the second lecture of [7].

Lemma 5.9. *For* $N \in \mathbb{N}$, *the* N*-parameter frozen percolation process is adapted to the filtration generated by the random variables* $\tau_v, v \in V$.

Recall the notation $\mathcal{C}_a(v; \lambda)$ from Definition 3.4. We prove the following proposition.

Proposition 3.5. *For all* $\lambda \in \mathbb{R}$ *and* $\varepsilon, K, \alpha > 0$, *there exist* $\theta = \theta(\lambda, \alpha, \varepsilon, K) > 0$ *and* $N_0 = N_0(\lambda, \alpha, \varepsilon, K)$ *such that*

$$\mathbb{P}_N(\exists v \in B(KN) \text{ s.t. } diam(\mathcal{C}_a(v; \lambda)) \in ((\alpha - \theta)N, (\alpha + \theta)N)) < \varepsilon$$

for $N \geq N_0$.

Proof of Proposition 3.5. Due to the length of the proof, we first give an outline.

For simplicity, we only give a sketch which shows that we can choose $\theta \in \left(0, \frac{1 \wedge \alpha}{2}\right)$ such that

$$\mathbb{P}_N(diam(\mathcal{C}_a(\lambda)) \in ((\alpha - \theta)N, (\alpha + \theta)N)) < \varepsilon \tag{5.11}$$

for large N.

Let us denote by $\tilde{x}, \tilde{y}$ a pair of sites in the active cluster of the origin for which $d(\tilde{x}, \tilde{y}) = diam(\mathcal{C}_a(\lambda))$. We consider the case where $\tilde{x}$ is one of the lowest and $\tilde{y}$ is one of the highest vertices of the active cluster. The other case where the diameter is achieved as a distance between a leftmost and rightmost vertex can be treated in a similar way. Let x (y) denote a vertex which is a neighbour of $\tilde{x}$ ($\tilde{y}$), and lies below (above) it. Note that x and y are closed frozen vertices at time $p_\lambda(N)$.

In Step 1 we apply Observation 3.2 and Lemma 2.27 to set λ_0 so that with probability close to 1, there are no frozen clusters at time $p_{\lambda_0}(N)$ in $B((\alpha + 2)N)$. Then we investigate the configuration close to x. In Step 2, we show that with probability close to 1, there is a unique frozen cluster F close to x. By Step 1, we can assume that it froze at time $p_{\lambda_F}(N)$ for $\lambda_F \in [\lambda_0, \lambda]$. In Step 4, we show that with probability close to 1, there is a graph $\mathcal{R} \subseteq \mathbb{T}$ such that its boundary consists of a $p_{\lambda_F}(N)$-closed arc, denoted by r_c, and a $p_{\lambda_F}(N)$-open arc. In Step 3,5 and 6 we show that with probability close to 1, we can impose some extra conditions on $\mathcal{R}$ and r_c and on the configuration in $\mathcal{R}$. We get a pair $(\mathcal{R}, r_c)$ with the following properties:

- $\partial\mathcal{R}$ is a certain outermost circuit, which is measurable with respect to the τ-values in $\mathbb{T} \setminus \mathcal{R}$, (Step 4)

- x is one of the the lowest vertices of $\mathcal{R}$ with two non-touching $p_{\lambda_F}(N)$-closed arms in $\mathcal{R}$ to r_c, (Step 4)
- no matter how we change the τ values in $\mathcal{R}$, the N-parameter frozen percolation outside $\mathcal{R}$ does not change up to time $p_\lambda(N)$, (Step 3)
- satisfies a technical condition (Step 5)
- $y \in \mathbb{T} \backslash cl(\mathcal{R})$ (Step 4).

Let us condition on the τ-values in $\mathbb{T} \setminus \mathcal{R}$. By the first and second property of $(\mathcal{R}, r_c)$ we get that in the N-parameter frozen percolation process the distribution of open and closed sites in $\mathcal{R}$ at time $p_{\lambda_F}(N)$ is the same as the distribution of ordinary site percolation in $\mathcal{R}$ with parameter $p_{\lambda_F}(N)$. This combined with $y \in \mathbb{T} \setminus \mathcal{R}$ allows us to decouple the locations of x and y. Since $d(\tilde{x}, \tilde{y}) = diam(\mathcal{C}_a(\lambda))$, to prove (5.11), it is enough to show that the first coordinate of x is not concentrated when we condition on the configuration in $\mathbb{T} \setminus \mathcal{R}$. We would like to use Corollary 5.8 for the pair $(\mathcal{R}, r_c)$. Unfortunately, this pair $(\mathcal{R}, r_c)$ might not satisfy all the conditions of Definition 5.5. To solve this problem we use the technical condition of Step 5 and we construct a pair $\left(\tilde{\mathcal{R}}, \tilde{r}_c\right)$ in Step 6 such that

- $\tilde{\mathcal{R}} \subset \mathcal{R}$,
- a translated version of $\left(\tilde{\mathcal{R}}, \tilde{r}_c\right)$ is $(\alpha_3 N, \alpha_2 N)$-regular as of Definition 5.5 for some $\alpha_2, \alpha_3 > 0$, and
- x is one of the the lowest vertices of $\tilde{\mathcal{R}}$ with two non-touching $p_{\lambda_F}(N)$-closed arms in $\tilde{\mathcal{R}}$ to $\tilde{r}_c$.

We apply Corollary 5.8 to $\left(\tilde{\mathcal{R}}, \tilde{r}_c\right)$ and get the required non-concentration result and finish the proof of Proposition 3.5. We make this argument precise in Step 7.

Remark. The structure of the proofs in Step 2-6 is an arm event hunting procedure: we take a some small neighbourhood of x. We deduce that if the required condition is violated, then we certain mixed near-critical arm events or crossing events of thin parallelograms occur. These events have upper bounds with exponents strictly larger than 2. Hence as we decrease the size of the neighbourhood, the probability of the occurrence of any of these events in the parallelogram $B((2\alpha + K + 2)N)$ goes to 0. We set the size of the neighbourhood such that the probability of the event where the condition of the step is not satisfied is as small as required. This finishes the proof of the step.

Let us turn to the precise proof.

Step 1. *We set* λ_0 *such that with probability close to* 1, *at time* $p_{\lambda_0}(N)$, *none of the open clusters intersecting* $B((2\alpha + K + 2)N)$ *are frozen.*

By Lemma 2.27 we choose $\lambda_0 = \lambda_0(\alpha, \varepsilon, K)$ and $N_0 = N_0(\alpha, \varepsilon, K)$ such that the event

$$E_0 := \mathcal{N}_c(\lambda_0, 1/24, 2\alpha + K + 4, N)$$

has probability at least $1 - \varepsilon/20$ for $N \geq N_0$. Then by Observation 3.2 we have that none of open clusters intersecting $B((2\alpha + K)N)$ are frozen. In particular, if a vertex $v \in B((2\alpha + K)N)$ is closed at time $p_\lambda(N)$, then it is $p_{\lambda_0}(N)$-closed. Moreover, if $v \in B((2\alpha + K + 2)N)$ is open at time $p_\lambda(N)$, then it is $p_\lambda(N)$-closed. This finishes Step 1.

Let $\theta \in \left(0, \frac{1 \wedge \alpha}{2}\right)$. For $i = 1, 2$, let $BA^i = BA^i(\theta)$ denote the set of vertices $v \in B(KN)$ such that there are $\tilde{x}(v) = (\tilde{x}_1(v), \tilde{x}_2(v))$, $\tilde{y}(v) = (\tilde{y}_1(v), \tilde{y}_2(v)) \in \mathcal{C}_a(v; \lambda)$ such that

$$\tilde{y}_i(v) - \tilde{x}_i(v) = d(\tilde{x}(v), \tilde{y}(v)) = diam(\mathcal{C}_a(v; \lambda)) \in ((\alpha - \theta)N, (\alpha + \theta)N). \tag{5.12}$$

Note that

$$\{\exists v \in B(KN) \text{ s.t. } diam(\mathcal{C}_a(v; \lambda)) \in ((\alpha - \theta)N, (\alpha + \theta)N)\} = \left\{BA^1 \cup BA^2 \neq \emptyset\right\}. \tag{5.13}$$

Let $u \in BA^2$. In the following we define quantities which depend on the value of u. In notation we indicate the dependence on u in the first appearance of these quantities, or when we want to emphasize this dependence. For each $u \in BA^2$ we fix a pair $(\tilde{x}, \tilde{y}) = (\tilde{x}, \tilde{y})(u)$ which satisfies (5.12). It can happen that there are more than one candidates for $\tilde{x}$ or $\tilde{y}$. In this case we choose one of them in some deterministic way. (E.g we can set $\tilde{x}$ ($\tilde{y}$) as the leftmost vertex among the candidates.) Let $x = x(u)$ ($y(u)$) denote a neighbour of $\tilde{x}$ ($\tilde{y}$) below $\tilde{x}$ (above $\tilde{y}$). The active cluster $\mathcal{C}_a(u; \lambda)$ lies between the horizontal lines passing through x and y denoted by e_x and e_y. Since



Figure 3: The closed boundary of $\mathcal{C}_a(\lambda)$ give rise to the closed arms c_L and c_R from x to $\partial B(x;\alpha N/2)$. The frozen vertex neighbouring x provides the arm o_B.

$\theta < \alpha/2$, the outer boundary of $\mathcal{C}_a(u;\lambda)$ provides two non-touching closed half plane arms in $x + \mathbb{Z} \boxtimes [0,\infty)$ to distance $\alpha N/2$ starting from x. Since $\partial\mathcal{C}_a(u;\lambda) \subset B((2\alpha+K+2)N)$, by Step 1, on the event E_0 these arms are $p_{\lambda_0}(N)$-closed. We denote the one on the left (right) hand side by $c_L = c_L(u)$ ($c_R = c_r(u)$). Apart from their common starting point, c_L and c_R do not even touch, since any active path connecting $\tilde{x}$ to $\tilde{y}$ separates them. Since x is a closed frozen vertex, there is at least one open frozen neighbour of x. From this vertex there is a $p_\lambda(N)$-open arm $o_B = o_B(u)$ to distance at least $N/2$. See Figure 3 for more details.

Let $\beta, \beta' \in (0,1)$ with $\beta < \beta'$. Recall the definition of the events $\mathcal{NA}(\beta,\beta') := \mathcal{NA}(\beta,\beta',\lambda,\lambda_0,2\alpha+K+2,N)$ and $\mathcal{NC}(\beta,\beta') := \mathcal{NC}(\beta,\beta',\lambda,\lambda_0,2\alpha+K+2,N)$ from Corollary 2.13 and 2.22. In the following we introduce the constants $\alpha_i > 0$ for $i = 1,2,3$ such that $\alpha_i/\alpha_{i+1} \gg 1$. Let $\alpha_3 \in \left(0, \frac{\alpha\wedge 1}{2}\right)$. Let $z = z(u) \in V$ such that $x = x(u) \in [-\alpha_3 N, \alpha_3 N] \boxtimes (-\alpha_3 N, \alpha_3 N]$. Note that $z \in B\left(\left\lceil \frac{2\alpha+K+2}{\alpha_3} \right\rceil\right)$. We define $B_3 = B_3(u) := B(\lfloor \alpha_3 N \rfloor z; \alpha_3 N)$.

Step 2. *We show that with probability close to* 1, *there is only one frozen cluster close to* $x = x(u)$ *for all* $u \in BA^2$.

Let $\alpha_1 \in \left(0, \frac{\alpha\wedge 1}{2}\right)$, $B_1 = B_1(u) := B(\lfloor \alpha_3 N \rfloor z; \alpha_1 N)$ and $A_1 = A_1(u) := A\left(\lfloor \alpha_3 N \rfloor z; \alpha_1 N, \frac{\alpha\wedge 1}{2} N\right)$. Suppose that there are at least two different frozen clusters in B_1. On the event E_0 we find $5,2$ mixed near critical arms in A_1 : the two $p_{\lambda_0}(N)$-closed arms c_L and c_R, the two $p_\lambda(N)$-open arms from the two frozen clusters, and a $p_{\lambda_0}(N)$-closed arm separating them. Let $E_1 := \mathcal{NA}\left(\alpha_1, \frac{\alpha\wedge 1}{2}\right)$. Hence we get:

Claim 5.10. On the event $E_0 \cap E_1$, $\forall u \in BA^2$, there is a unique frozen cluster denoted by $F = F(u)$ which intersects $B_1(u)$. Let $\lambda_F = \lambda_F(u) \in [\lambda_0, \lambda]$ such that F froze at $p_{\lambda_F}(N)$. On $E_0 \cap E_1$, a vertex in $B_1(u)$ is open in the N-parameter frozen percolation process at time $p_{\lambda_F}(N)$ if and only if it is $p_{\lambda_F}(N)$-open.

In the following two steps we write open (closed) for $p_{\lambda_F}(N)$-open ($p_{\lambda_F}(N)$-closed) if it is not stated otherwise. We finish Step 2 by applying Corollary 2.13 and we set α_1 such that

$$\mathbb{P}(E_1) \geq 1 - \varepsilon/20 \tag{5.14}$$

for $N \geq N_1(\varepsilon, \lambda_0, \lambda, \alpha, K)$.

Step 3. *We say that a circuit is $p_{\lambda_F}(N)$-open-closed, or simply open-closed, if it consists of a $p_{\lambda_F}(N)$-open and a $p_{\lambda_F}(N)$-closed arc. Suppose that there is a $p_{\lambda_F}(N)$-open-closed circuit close to and around x. We show that with probability close to* 1*, no matter how we change the τ values inside this circuit, the N-parameter frozen percolation process does not change till time $p_\lambda(N)$ outside of the circuit.*

Let $\alpha_2 \in \left(0, \alpha_1 \wedge \frac{1}{4}\right)$, and $\beta_2 \in (\alpha_2, \alpha_1)$ be some intermediate scale. We define $B_2 = B_2(u) := B(\lfloor \alpha_3 N \rfloor z; \alpha_2 N)$, $B_2' = B_2'(u) := B(\lfloor \alpha_3 N \rfloor z; \beta_2 N)$, $A_2 = A_2(u) := A(\lfloor \alpha_3 N \rfloor z; \alpha_2 N, \alpha_1 N)$ and $A_2' = A_2'(u) := A(\lfloor \alpha_3 N \rfloor z; \beta_2 N, \alpha_1 N)$. Let $BL = BL(u)$ denote the set of bordering lines of $F \setminus B_2'$, that is the top- and bottom-most horizontal, left- and rightmost vertical lines which intersect $F \setminus B_2'$. We rule out the case where there is a line in BL which intersects B_2' in the following technical claim.

Claim 5.11. Let

$$E_2' = \mathcal{NA}(2\beta_2, \alpha_1 - 2\beta_2) \cap \mathcal{NC}(2\beta_2, 2\alpha_1). \tag{5.15}$$

Then

$$E_0 \cap E_1 \cap E_2' \subset E_0 \cap E_1 \cap \left\{\forall u \in BA^2, \forall e \in BL(u) \text{ we have } e \cap (F \setminus B_2') = \emptyset\right\}.$$

Proof of Claim 5.11. Let $u \in BA^2$. When the bottom-most line of $F \setminus B_2'$ intersects B_2', then $F \subseteq (\mathbb{Z} \boxtimes [-\beta_2 N, \infty)) + \lfloor \alpha_3 N \rfloor z$. We see 4 half plane arms: c_L, c_R give two closed and o_B gives an open arm, a fourth closed half plane arm separates F from the line $\mathbb{Z} \boxtimes \{\lfloor \beta_2 N \rfloor\} + \lfloor \alpha_3 N \rfloor z$. Hence $\mathcal{NA}^c(2\beta_2, \alpha_1 - 2\beta_2)$ occurs.

If the topmost line of $F \setminus B_2'$ intersects B_2', then the closed arms c_L and c_R stay in the parallelogram

$$[-\alpha_1 N, \alpha_1 N] \boxtimes [-\beta_2 N, \beta_2 N] + \lfloor \alpha_3 N \rfloor z.$$

Hence we get a closed horizontal crossing of it, that is the event $\mathcal{NC}^c(2\beta_2, 2\alpha_1)$ occurs.

When a leftmost bordering line of $F \setminus B_2'$ intersects B_2', then we find that the arms in $A(\lfloor \alpha_3 N \rfloor z; \beta_2 N, \alpha_1 N)$ induced by c_L, c_R and o_B stay in half plane

$$[-2\beta_2 N, \infty) \times \mathbb{R} + \lfloor \alpha_3 N \rfloor z. \tag{5.16}$$

The frozen cluster F is separated from the line $\{-2\beta_2 N\} \times \mathbb{R} + \lfloor \alpha_3 N \rfloor z$. This provides an additional closed arm in the half plane (5.16), which together the arms induced by c_L, c_R and o_B give 4 half plane arms, hence the event $\mathcal{NA}^c(\beta_2, \alpha_1 - 2\beta_2)$ occurs.

The case when the rightmost bordering line of $F \setminus B_2'$ intersects B_2' can be treated similarly.

With the notation (5.15) we get that on the event $E_0 \cap E_1 \cap E_2'$, none of the lines of BL intersect B_2', which finishes the proof of Claim 5.11. ☐

Now we proceed with Step 3. Let $u \in BA^2$. Suppose that there is an open-closed circuit $OC = OC(u)$ around x in B_2. Let $R = R(u)$ denote the the union of the finite connected components of $\mathbb{T} \setminus OC$. Let us change the τ values of the vertices in R in some arbitrary non-degenerate way (that is, the new τ values are all different), but keep the original values outside R. Let us run the N-parameter frozen percolation dynamics for this modified set of τ values.

First we consider the case where this new process differs from the old one outside of R at some time $t \in [0, p_{\lambda_F}(N)]$. The closed arc of OC stays closed till time $p_{\lambda_F}(N)$ in both processes. Hence it acts as a barrier for the effect of τ values in R. The open arc of OC is a subset of F. By Claim 5.11 on the event $E_0 \cap E_1 \cap E_2'$ if these two processes differ outside R, then in the new one a frozen cluster F' emerged before time $p_{\lambda_F}(N)$. Since the two processes differ outside R, we have $F' \setminus R \neq F \setminus R$. On the event E_0 there are no $p_{\lambda_0}(N)$-open clusters with diameter at least $N/4$ intersecting $B((2\alpha + K + 2)N)$. Since $\alpha_2 < 1/4$ we get that F' froze in the new process at time $p_{\lambda_{F'}}(N)$ with $\lambda_{F'} \in [\lambda_0, \lambda_F]$. Let BL' denote the set of bordering lines of $F' \setminus B_2'$. Since $\lambda_{F'} \in [\lambda_0, \lambda_F]$, the arguments of the proof of Claim 5.11 applied to the new process give that on the event $E_0 \cap E_1 \cap E_2'$ none of the lines of BL' intersect B_2'. Hence $F' \setminus R$ has two connected components F_1' and F_2' such that $\mathit{diam}(F_i') < N$ for $i \in \{1, 2\}$, but $\mathit{diam}(F_1' \cup F_2') \geq N$. Since $R \subset B_2$, each of F_1', F_2' contains a $p_{\lambda_{F'}}(N)$-open arm in the annulus $A_2'' = A_2''(u) := A(\lfloor \alpha_3 N \rfloor z; \alpha_2 N, \beta_2 N)$. When for some $i \in \{1, 2\}$ F_i' lies above c_L and c_R, then we get a 4, 3 near critical arm event: the closed arms induced by c_L, c_R and the open arm induced by F_i' stay above e_x, and o_B provides the fourth arm in A_2''. Hence $\mathcal{NA}^c(\alpha_2, \beta_2)$ occurs. If both of F_1', F_2' lie below c_L and c_R then we get a 5, 2 near critical mixed arm event in A_2'' : c_L, c_R induce closed half plane arms in A_2''. F_1', F_2' induce two open arms. Since F_1' and F_2' are different connected components of $F' \setminus R$, there is a fifth, $p_{\lambda_{F'}}(N)$-closed, arm separating F_1' and F_2' in A_2''. Hence $\mathcal{NA}^c(\alpha_2, \beta_2)$ occurs. Let $E_2 = E_2' \cap \mathcal{NA}(\alpha_2, \beta_2)$.

Let us consider the other case where the new and the old process differ from each other outside of R at some time $t \in (p_{\lambda_F}(N), p_\lambda(N)]$, but they agree outside of R till time $p_{\lambda_F}(N)$. Since till time $p_{\lambda_F}(N)$ the two processes coincide outside of R, then a frozen cluster F' is formed at time $p_{\lambda_F}(N)$ in the new process. Moreover, $F' \setminus R = F \setminus R$.

However, the two processes differ at some time $t \in (p_{\lambda_F}(N), p_\lambda(N)]$, hence an additional frozen cluster F'' has to emerge in this time period using some of the vertices in R. This induces the $5, 2$ near critical mixed arm event of Step 1. Hence we proved the following claim.

Claim 5.12. On the event $E_0 \cap E_1 \cap E_2$, we have that $\forall u \in BA^2$, if there is a $p_{\lambda_F}(N)$-open-closed circuit around $x = x(u)$ in $B_2(u)$ then no matter how we change the τ values inside this circuit, the frozen percolation process outside it does not change till time $p_\lambda(N)$.

We finish Step 3 by applying Corollary 2.13 and 2.24: we fix the value of β_2', β_2 and α_2 such that

$$\mathbb{P}(E_2) \geq 1 - \varepsilon/20 \tag{5.17}$$

for $N \geq N_2(\varepsilon, \lambda_0, \lambda, \alpha, K)$.

Step 4. *We show that with probability close to* 1, *there is a* $p_{\lambda_F}(N)$*-open-closed circuit around* x, *such that the location where its colour changes in the circuit is 'far' above* x.

Let $u \in BA^2$. Let $\alpha_3 \in (0, \alpha_2)$, $B_3 = B_3(u) := B(\lfloor \alpha_3 N \rfloor z; \alpha_3 N)$ and $A_3 = A_3(u) := A(\lfloor \alpha_3 N \rfloor z; \alpha_3 N, \alpha_2 N)$. Let $\delta_3 \in (\alpha_3, \alpha_2)$ be an intermediate scale. We cut the annulus A_3 into three subannuli using two other intermediate scales β_3, β_3' with $\alpha_3 < \delta_3 < \beta_3 < \beta_3' < \alpha_2$:

$$\begin{aligned} A_{3,0} &= A_{3,0}(u) := A(\lfloor \alpha_3 N \rfloor z; \alpha_3 N, \beta_3 N), \\ A_{3,1} &= A_{3,1}(u) := A(\lfloor \alpha_3 N \rfloor z; \beta_3 N, \beta_3' N), \\ A_{3,2} &= A_{3,2}(u) := A(\lfloor \alpha_3 N \rfloor z; \beta_3' N, \alpha_2 N). \end{aligned}$$

Let $\bar{c}_L$ ($\bar{c}_R$) denote the closed arm induced by c_L (c_R) in $A_{3,1}$.

If c_L and c_R are not connected by a closed path in $A_{3,0} \cap \mathcal{C}_a(u; \lambda)$, then there is a open arm separating them. Hence we see a near critical $4, 3$ arm event: c_L, c_R and the separating open arm induce disjoint half plane arms in $A_{3,0}$, and the fourth arm in $A_{3,0}$ is induced by o_B. Thus the event $\mathcal{NA}^c(\alpha_3, \beta_3)$ occurs.

If $\bar{c}_L \subseteq [-\beta_3' N, -\beta_3 N] \boxtimes [-\alpha_3 N, \delta_3 N]$ or $\bar{c}_R \in [\beta_3 N, \beta_3' N] \boxtimes [-\alpha_3 N, \delta_3 N]$, then we find a closed horizontal crossing in a narrow parallelogram. Hence the event $\mathcal{NC}^c(2\delta_3, \beta_3' - \beta_3)$ occurs.

In the following we assume that both $\bar{c}_L$ and $\bar{c}_R$ leave the corresponding parallelograms. Let w_L (w_R) be an open frozen vertex neighbouring a vertex of $\bar{c}_L$ ($\bar{c}_R$) which is outside of the aforementioned parallelogram.

Suppose that there is no open arc in A_3 connecting w_L to o_B. Since w_L is open frozen at time $p_{\lambda_F}(N)$, it has a $p_{\lambda_F}(N)$-open path path to distance $N/2$. Let o_L denote the part of this path till the first time it exists A_3. Note that o_L and o_B are disjoint, and they are not connected by open path inside A_3. We have two cases depending on where o_B leaves A_3.

When it leaves A_3 by exiting its outer parallelogram, than we get a $5, 2$ near critical arm event in $A_{3,2}$: two half plane closed arms induced by c_L and c_R, two open arms induced by o_L and o_B an extra closed arm separates o_L and o_B in $A_{3,2}$. Hence the event $\mathcal{NA}^c(\beta_3', \alpha_2)$ occurs. See Figure 4.

When o_L leaves A_3 by entering its inner parallelogram, then we get a similar $5, 2$ arm event in $A_{3,0}$. Thus $\mathcal{NA}^c(\alpha_3, \beta_3)$ happens. In a similar way we can show that when w_R is not connected to o_B in $A_{3,x}$, then $\mathcal{NA}^c(\beta_3', \alpha_2) \cup \mathcal{NA}^c(\alpha_3, \beta_3)$ occurs.

See Figure 5. Note that $w_L, w_R \in (\mathbb{Z} \boxtimes [\delta_2 N, \alpha_2 N]) + \lfloor \alpha_5 N \rfloor z$. With the notation

$$E_3 = \mathcal{NC}(\alpha_3 + \delta_3, \beta_3' - \beta_3) \cap \mathcal{NA}(\alpha_3, \beta_3) \cap \mathcal{NA}(\beta_3', \alpha_2) \cap \mathcal{NA}(\alpha_3, \beta_3)$$

we proved the following claim.

Claim 5.13. We have that on the event $E_0 \cap E_1 \cap E_2 \cap E_3$, $\forall u \in BA^2$ there is a $p_{\lambda_{F(u)}}(N)$-open-closed circuit in $A_3(u)$ such that locations where the colour changes in the circuit is $(\mathbb{Z} \boxtimes [\delta_3 N, \alpha_2 N]) + \lfloor \alpha_3 N \rfloor z$.

We finish Step 4 by choosing the values of β_3, β_3' and δ_3. The probability of E_3 is an increasing function of α_3 for $\beta_3, \beta_3', \delta_3$ fixed. By Corollary 2.13 and 2.24 we choose the value of $\beta_3, \beta_3', \delta_3, \alpha_3$ such that the probability of the event E_3 is at least $1 - \varepsilon/20$. We only fix $\beta_3, \beta_3', \delta_3$ and require α_3 to be small but unspecified so that

$$\mathbb{P}(E_3) \geq 1 - \varepsilon/20 \tag{5.18}$$

for $N \geq N_3(\varepsilon, \alpha_3, \lambda_0, \lambda, \alpha, K)$. We choose the value of α_3 in Step 6.

Before Step 5, let us summarize what we have proved up to now. Let $u \in BA^2$, and suppose that the event $E_0 \cap E_1 \cap E_2 \cap E_3(\alpha_3)$ holds. Consider the circuit OC which we constructed in Step 4. It has a special property: as we walk from the outside of $B_2 = B(\lfloor \alpha_3 N \rfloor z; \alpha_2 N)$ on one of the closed arms c_L and c_R towards x, we hit the



Figure 4: The closed arm c_{LB} separates o_L and o_B in $A_{3,2,x}$. Hence $c_L, c_R, o_B, c_{LB}, o_L$ give $5, 2$ near critical mixed arms.



Figure 5: The circuit around $B_{3,x}$ consists of the open arc drawn with continuous line, subpaths of c_L and c_R and the closed arc in $A_{3,0,x}$.

closed part of OC at its endpoints for the first time. See Figure 5 for more details. Let $\mathcal{OC}$ denote the outermost open-closed circuit in A_3 satisfying the conditions appearing Claim 5.13 and this special property. Let $\mathcal{R}$ denote the union of finite connected components of $\mathbb{T} \setminus \mathcal{OC}$. Let r_o and r_c denote the open and closed parts of $\mathcal{OC} = \partial\mathcal{R}$. The pair $(\mathcal{R}, r_c)$ and the configuration in $\mathbb{T}\backslash\mathcal{R}$ satisfies the following conditions:

1. $\mathcal{R}$ is an induced subgraph of $\mathbb{T}$ such that $cl(\mathcal{R})$ is connected, (by Claim 5.13)
2. $B(\lfloor\alpha_3 N\rfloor z; \alpha_3 N) = B_3 \subseteq \mathcal{R} \subseteq B_2 = B(\lfloor\alpha_3 N\rfloor z; \alpha_2 N)$ (by Claim 5.13)
3. $\partial\mathcal{R}$ is disjoint union of non-empty self avoiding paths r_c and r_o, which are oriented such that $\mathcal{R}$ lies on the right when we walk along them, (by Claim 5.13)
4. $r_c \subseteq [-\alpha_2 N, \alpha_2 N] \boxtimes [-\alpha_3 N, \alpha_2 N] + \lfloor\alpha_5 N\rfloor z$, (by the proof of Claim 5.13)
5. the endpoints of r_c denoted by s_L and s_R lie in the parallelogram $[-\alpha_3 N, \alpha_3 N] \boxtimes [\delta_3 N, \alpha_3 N] + \lfloor\alpha_3 N\rfloor z$, (by Claim 5.13)
6. when we walk along c_L (c_R) towards x, we hit $\partial\mathcal{R}$ first at vertex s_L (s_R), (by the proof of Claim 5.13)
7. for every vertex $v \in r_o$, there is a closed path in $B_2 \setminus \mathcal{R}$ to ∂B_2, ($\mathcal{OC}$ is outermost)
8. for every vertex $v \in r_c$, there is an open path in $B_2 \setminus \mathcal{R}$ to ∂B_2 or to $(c_L \cup c_R) \setminus cl(\mathcal{R})$. ($\mathcal{OC}$ is outermost)

Note that the first three conditions coincide with the first three conditions for the pair $(\mathcal{R} - \lfloor\alpha_3 N\rfloor z, r_c - \lfloor\alpha_3 N\rfloor z)$ being $(\alpha_3 N, \alpha_2 N)$-outer-regular of Definition 5.5. We add an extra condition in the next step.

Note that the vertex x has two non-touching closed arms to r_c. Moreover, by Condition 6, x is one of the lowest vertices in $\mathcal{R}$ with this property. With the notation of Definition 5.1 we have that $x \in \mathcal{L}(\mathcal{R}, r_c)$ in the N-parameter frozen percolation process at time $p_{\lambda_F}(N)$.

Step 5. Let $u \in BA^2$. *Suppose that the event* $E_0 \cap E_1 \cap E_2 \cap E_3$ *holds.* Let $\mathcal{W} = \mathcal{W}(u)$ denote the connected components of $\mathcal{R} \cap (\mathbb{Z} \boxtimes [-\lfloor\alpha_3 N\rfloor + 1, \lfloor 5\alpha_3 N\rfloor - 1])$. *Let S_M denote the unique element of $\mathcal{W}$ which contains B_3 as a subset. We show that with probability close to* 1, $\partial S_M \cap r_c = \emptyset$.

We define $e_T = e_T(u) := (\mathbb{Z} \boxtimes \{\lfloor 5\alpha_3 N\rfloor + 1\}) + \lfloor\alpha_3 N\rfloor z$ and $e_B = e_B(u) := (\mathbb{Z} \boxtimes \{-\lfloor\alpha_3 N\rfloor - 1\}) + \lfloor\alpha_3 N\rfloor z$. Suppose that $\partial S_M \cap r_c \neq \emptyset$, let $w \in \partial S_M \cap r_c \cap e_T$. Consider the parallelogram $\bar{B} = B(s; \delta_3 N/2)$. Let w_L and w_R denote the vertices of r_c where we exit $\bar{B}$ the first time as we walk on r_c starting from s towards s_L and s_R. The part of r_c between w_L and w_R cuts $\bar{B}$ into two pieces. Let $\bar{B}_I$ ($\bar{B}_E$) denote the part which is on the right (left) hand side of r_c when we walk from w_L to w_R. Let $\bar{A}_I = \bar{B}_I \setminus B(s; 6\alpha_3 N)$ and $\bar{A}_E = \bar{B}_E \setminus B(s; 6\alpha_3 N)$. By Condition 8 above $\bar{A}_E$ contains an open arm. We claim that $\bar{A}_I$ also contains an open arm. Suppose the contrary. Then there must be a closed non self-touching arc in $\bar{A}_I$ preventing the occurrence of the open arm. Note that this arc is contained in $\mathcal{R}$. Then the lowest vertex of this arc has two disjoint $p_{\lambda_F}(N)$-closed arms to r_c, and it lies lower than $x \in B := B(\lfloor\alpha_3 N\rfloor z; \alpha_3 N)$. This contradicts $x \in \mathcal{L}(\mathcal{R}, r_c)$ which was shown in the lines before Step 4. See Figure 6. Hence $\bar{A}_I$ has an open arm, which together the open arm of $\bar{A}_E$ and the two closed arms of w provide a 4, 3 near critical mixed arm event.Hence the event $E_4^c = \mathcal{NA}^c(6\alpha_3, \delta_3/2)$ occurs. Thus we arrive to the following claim which finishes Step 5.

Claim 5.14. On the event $E_0 \cap E_1 \cap E_2 \cap E_3 \cap E_4$, we have $\partial S_M \cap r_c = \emptyset$.

Step 6. *Recall Definition 5.5. We show that with probability close to* 1, *we can cut down some parts of $\mathcal{R}$ and get a pair $\tilde{\mathcal{R}}$ and $\tilde{r}_c$ such that the pair $\left(\tilde{\mathcal{R}} - \lfloor\alpha_3 N\rfloor z, \tilde{r}_c - \lfloor\alpha_3 N\rfloor z\right)$ is $(\alpha_3 N, \alpha_2 N)$-regular and*

$$\mathcal{L}(\mathcal{R}, r_c) \cap B = \mathcal{L}\left(\tilde{\mathcal{R}}, \tilde{r}_c\right) \cap B.$$

Let $u \in BA^2$. Suppose that the event $E_0 \cap E_1 \cap E_2 \cap E_3 \cap E_4$ occurs. Let $\tilde{\mathcal{R}} = \tilde{\mathcal{R}}(u)$ be the connected component of S_M in $\mathcal{R} \setminus \bigcup_{S \in \mathcal{W}: \partial S \cap r_c \neq \emptyset} cl(S)$ and $\tilde{r}_c = \partial\tilde{\mathcal{R}} \setminus r_o$. The conditions before Step 5 and Claim 5.14 gives that the pair $\left(\tilde{\mathcal{R}} - \lfloor\alpha_3 N\rfloor z, \tilde{r}_c - \lfloor\alpha_3 N\rfloor z\right)$ is $(\alpha_3 N, \alpha_2 N)$-regular.

For $R \subset \mathbb{T}$ and $r \subset \partial R$ let $\mathcal{TA}(R, r)$ denote the set of closed vertices $v \in R$ such that v has two non-touching closed arms in R to r. Let M denote the connected component of S_M in $R \setminus e_T$. We show the following:



Figure 6: The grey area represents $\bar{A}_I$. If there is no open arm in $\bar{A}_I$ then we there is a closed arc in $\bar{A}_I$. This contradicts with x being one of the lowest vertices of $\mathcal{C}_a(\lambda)$.

Claim 5.15. Let

$$E_5 := \mathcal{NA}(6\alpha_3, \beta_4) \cup \mathcal{NA}(\beta_4, \delta_3/2). \tag{5.19}$$

On the event $\bigcap_{i=0}^5 E_i$ $\forall u \in BA^1$, the pair $\left(\tilde{\mathcal{R}} - \lfloor \alpha_3 N \rfloor z, \tilde{r}_c - \lfloor \alpha_3 N \rfloor z\right)$ is $(\alpha_3 N, \alpha_2 N)$-regular, and

$$\mathcal{TA}(\mathcal{R}, r_c) \cap M = \mathcal{TA}\left(\tilde{\mathcal{R}}, \tilde{r}_c\right) \cap M.$$

In particular,

$$\mathcal{L}(\mathcal{R}, r_c) \cap B = \mathcal{L}\left(\tilde{\mathcal{R}}, \tilde{r}_c\right) \cap B.$$

Proof of Claim 5.15. From the definition of $\left(\tilde{\mathcal{R}}, \tilde{r}_c\right)$ it follows that $(\mathcal{TA}(\mathcal{R}, r_c) \cap M) \subset \left(\mathcal{TA}\left(\tilde{\mathcal{R}}, \tilde{r}_c\right) \cap M\right)$. Suppose that $\exists v \in \left(\mathcal{TA}(\mathcal{R}, r_c) \setminus \mathcal{TA}\left(\tilde{\mathcal{R}}, \tilde{r}_c\right)\right) \cap M$. Let c_v^1 and c_v^2 denote two non-touching closed arms starting from v and ending at $v^1 \in \tilde{r}_c$ and $v^2 \in \tilde{r}_c$ respectively. Since $v \in \mathcal{TA}(\mathcal{R}, r_c) \setminus \mathcal{TA}\left(\tilde{\mathcal{R}}, \tilde{r}_c\right)$, we can assume that c_v^1 cannot be extended in such a way that it connects to r_c and this extension is disjoint from and does not touch c_v^2. Hence $v^1 \in \tilde{r}_c \setminus r_c$, and $v^1 \in e_T$. Let $S \in \mathcal{W}$ such that $v^1 \in \partial S$. Note that $\partial S \cap r_c \neq \emptyset$. Let s^1, s^2 denote the endpoints of the connected component of v^1 in $\partial S \cap e_T$. At least one of s^1 and s^2 is in r_c. Let $s^1 \in r_c$. Let $\beta_4 \in (6\alpha_3, \delta_3/2)$ be an intermediate scale. We divide the annulus $A\left(v^1; 6\alpha_3 N, \delta_3 N/2\right)$ into the annuli

$$A_{4,0} = A\left(v^1; 6\alpha_3 N, \beta_4 N\right),$$
$$A_{4,1} = A\left(v^1; \beta_4 N, \delta_3 N/2\right).$$

We have two cases. If $c_v^2 \cap A_{4,0} \neq \emptyset$, then we see 4 half plane arms in $A_{4,1}$: r_c provides two closed arms, and each of c_v^1 and c_v^2 gives one closed arm. Hence the event $\mathcal{NA}^c(6\alpha_3, \beta_4)$ occurs. If $c_v^2 \cap A_{4,0} = \emptyset$, we have 4 half plane arms in $A_{4,0}$: r_c provides two closed arms, c_v^1 an other closed arm, moreover, we get an open arm which separates c_v^1 from r_c. See Figure 7 more details. Hence the event $\mathcal{NA}^c(\beta_4, \delta_3/2)$ occurs. By (5.19) this finishes the proof of Claim 5.15. □



Figure 7: If $c_v^2 \cap A_{4,0} = \emptyset$, we see 4 half plane arms in $A_{4,0}$: the two closed induced by r_c, a closed arm c_v^1, and an open arm o_v which separates c_v^1 from r_c.

By Corollary 2.13 we set α_3 such that

$$\mathbb{P}\left(E_4 \cap E_5\right) \geq 1 - \varepsilon/20 \tag{5.20}$$

for $N \geq N_5\left(\varepsilon, \alpha_3, \lambda_0, \lambda, \alpha, K\right)$. Let

$$E = E_0 \cap E_1 \cap E_2 \cap E_3 \cap E_4 \cap E_5.$$

The combination of the lines in the beginning of Step 1, (5.14), (5.17), (5.18) and (5.20) gives that

$$\mathbb{P}\left(E\right) \geq 1 - \varepsilon/4 \tag{5.21}$$

for $N \geq \bigvee_{i=0}^{5} N_i$. This finishes Step 6.

Step 7. *We set $\theta > 0$ such that $\mathbb{P}_N\left(BA^2 \neq \emptyset\right) < \varepsilon/2$ for large N, and conclude the proof of Proposition 3.5.*

For $v \in V$, let

$$Z\left(v\right) := \left\{\exists u \in BA^2 \text{ such that } z\left(u\right) = v\right\}.$$

Hence

$$\left\{BA^2 \neq \emptyset\right\} = \bigcup_{v \in B\left(\left\lceil \frac{2\alpha+K+2}{\alpha_3}\right\rceil\right)} Z\left(v\right)$$

and

$$\mathbb{P}_N\left(BA^2 \neq \emptyset, E\right) \leq \sum_{v \in B\left(\left\lceil \frac{2\alpha+K+2}{\alpha_3}\right\rceil\right)} \mathbb{P}_N\left(Z\left(v\right) \cap E\right) \tag{5.22}$$

Note that on the event $Z\left(v\right) \cap E$, Claim 5.10 and the arguments above give that $\mathcal{C}_a\left(u;\lambda\right), F\left(u\right), \lambda_F\left(u\right), \mathcal{R}\left(u\right), r_c\left(u\right), \tilde{\mathcal{R}}\left(u\right)$ and $\tilde{r}_c\left(u\right)$ do not depend on the choice of $u \in BA^2$ as long as $z\left(u\right) = v$. Except for $\mathcal{C}_a\left(u,\lambda\right)$, we omit the argument u from the notation above.

We set $k := \lfloor 1/2\theta \rfloor$. Recall that $d\left(x,y\right) = d\left(\tilde{x},\tilde{y}\right) + \sqrt{3} = diam\left(\mathcal{C}_a\left(u;\lambda\right)\right) + \sqrt{3}$, and $diam\left(\mathcal{C}_a\left(u;\lambda\right)\right) \in \left(\left(\alpha-\theta\right)N, \left(\alpha+\theta\right)N\right)$. On the event $Z\left(v\right)$ there is a unique $l = l\left(y\right) \in [0, k-1] \cap \mathbb{Z}$ such that $x \in B_{l,k}$ where

$$B_{l,k} = B_{l,k}\left(v\right) := \left[-\alpha_3 N, \alpha_3 N\right] \boxtimes \left(\left(2\frac{l}{k} - 1\right)\alpha_3 N, \left(2\frac{l+1}{k} - 1\right)\alpha_3 N\right] + \lfloor \alpha_3 N \rfloor v.$$

Recall from the lines above Step 5 we have $x \in \mathcal{L}(\mathcal{R}, r_c)$. From Claim 5.15 we have $\mathcal{L}(\mathcal{R}, r_c) \cap B = \mathcal{L}\left(\tilde{\mathcal{R}}, \tilde{r}_c\right) \cap B$ where $B = B(\lfloor \alpha_3 N \rfloor v; \alpha_3 N)$. Hence on the event $Z(v) \cap E$, we have $\mathcal{L}\left(\tilde{\mathcal{R}}, \tilde{r}_c\right) \cap B_{l,k} \neq \emptyset$. Let (R, r) be a fixed pair. Hence

$$\mathbb{P}_N(Z(v), E, (\mathcal{R}, r_c) = (R, r)) = \mathbb{P}_N\left(Z(v), E, (\mathcal{R}, r_c) = (R, r), \mathcal{L}\left(\tilde{R}, \tilde{r}\right) \cap B_{l,k} \neq \emptyset \text{ at time } p_{\lambda_F}(N)\right) \tag{5.23}$$

where $\left(\tilde{R}, \tilde{r}\right)$ denotes the pair we get when we cut down some parts of R as in Step 6.

For $t \in [0,1]$ and $J \subset V$ let

$$\mathcal{F}_t(J) := \sigma(\{\tau_w < s\} \,|\, w \in J, s \in [0,1])$$

denote the σ-algebra generated by the τ values of the vertices in J up to time t.

Lemma 5.9 gives that the N-parameter frozen percolation process is adapted to the filtration $(\mathcal{F}_t(V))_{t\in[0,1]}$. Hence for all $u \in BA^2$, $\{(\mathcal{R}, r_c) = (R, r)\}$, l and λ_F are measurable with respect to $\mathcal{F}_{p_\lambda(N)}(V)$. By Claim 5.12 we have that on the event $Z(v) \cap E \cap \{(\mathcal{R}, r_c) = (R, r)\}$ the τ-values in R do not influence the frozen percolation process in $V \setminus R$ up to time $p_\lambda(N)$. Since $\partial\mathcal{R}$ is a certain outermost circuit, the event $\left\{(\mathcal{R}, r_c) = (R, r), l = \bar{l}, \lambda_F = \bar{\lambda}_F\right\}$ is measurable with respect to $\mathcal{F}_{p_\lambda(N)}(V \setminus R)$ on the event $Z(v) \cap E$. Hence

$$\begin{aligned}
&\mathbb{P}_N\left(Z(v), E, (\mathcal{R}, r_c) = (R, r), l = \bar{l}, \lambda_F = \bar{\lambda}_F, \mathcal{L}\left(\tilde{R}, \tilde{r}\right) \cap B_{l,k} \neq \emptyset \text{ at time } p_{\lambda_F}(N) \,\middle|\, \mathcal{F}_{p_\lambda(N)}(V \setminus R)\right) \\
&\quad = \mathbf{1}\left\{(\mathcal{R}, r_c) = (R, r), l = \bar{l}, \lambda_F = \bar{\lambda}_F\right\} \mathbb{P}_N\left(Z(v), E, \mathcal{L}\left(\tilde{R}, \tilde{r}\right) \cap B_{\bar{l},k} \neq \emptyset \text{ at time } p_{\bar{\lambda}_F}(N) \,\middle|\, \mathcal{F}_{p_\lambda(N)}(V \setminus R)\right),
\end{aligned} \tag{5.24}$$

for $\bar{l} \in [0, k-1] \cap \mathbb{Z}$ and $\bar{\lambda}_F \in [\lambda_0, \lambda]$. In (5.24) the definition of $\mathbf{1}\left\{(\mathcal{R}, r_c) = (R, r), l = \bar{l}, \lambda_F = \bar{\lambda}_F\right\}$ requires some care. We define it as follows. Recall that the N-parameter frozen percolation process is adapted to the filtration $(\mathcal{F}_t(V))_{t\in[0,1]}$. Hence we can assume that the sample space is $\Omega = [0,1]^V$. Let $\omega \in [0,1]^{V\setminus R}$. We set $\mathbf{1}\left\{(\mathcal{R}, r_c) = (R, r), l = \bar{l}, \lambda_F = \bar{\lambda}_F\right\}(\omega) = 1$ if and only if there is $\omega' \in [0,1]^V$ such that

$$\omega' \in Z(v) \cap E \cap \left\{(\mathcal{R}, r_c) = (R, r), l = \bar{l}, \lambda_F = \bar{\lambda}_F\right\}$$

and $\omega(v) = \omega'(v)$ for $v \in V \setminus R$.

From Step 6 we have $\tilde{R} \subset R$. Claim 5.15 shows that we can apply Corollary 5.8 in the following. We have

$$\begin{aligned}
&\mathbb{P}_N\left(Z(v), E, \mathcal{L}\left(\tilde{R}, \tilde{r}\right) \cap B_{\bar{l},k} \neq \emptyset \text{ at time } p_{\bar{\lambda}_F}(N) \,\middle|\, \mathcal{F}_{p_\lambda(N)}(V \setminus R)\right) \\
&\qquad \leq \mathbb{P}_N\left(\mathcal{L}\left(\tilde{R}, \tilde{r}\right) \cap B_{\bar{l},k} \neq \emptyset \text{ at time } p_{\bar{\lambda}_F}(N) \,\middle|\, \mathcal{F}_{p_\lambda(N)}(V \setminus R)\right) \\
&\qquad = \mathbb{P}_{p_{\bar{\lambda}_F}(N)}\left(\mathcal{L}\left(\tilde{R}, \tilde{r}\right) \cap B_{\bar{l},k} \neq \emptyset\right) \\
&\qquad \leq c_1 k^{-1}
\end{aligned} \tag{5.25}$$

for $N \geq N_6(\lambda_0, \lambda, \alpha_3, \alpha_2, k)$ with $c_1 = c_1(\lambda_0, \lambda, \alpha_3, \alpha_2)$ of Corollary 5.8. A combination of (5.25) and (5.24) gives that

$$\begin{aligned}
&\mathbb{P}_N\left(Z(v), E, (\mathcal{R}, r_c) = (R, r), l = \bar{l}, \lambda_F = \bar{\lambda}_F, \mathcal{L}\left(\tilde{R}, \tilde{r}\right) \cap B_{l,k} \neq \emptyset \text{ at time } p_{\lambda_F}(N) \,\middle|\, \mathcal{F}_{p_\lambda(N)}(V \setminus R)\right) \\
&\qquad \leq c_1 k^{-1} \mathbf{1}\left\{(\mathcal{R}, r_c) = (R, r), l = \bar{l}, \lambda_F = \bar{\lambda}_F\right\}
\end{aligned}$$

for $N \geq N_6$. Hence

$$\mathbb{P}_N(Z(v) \cap E) \leq c_1 k^{-1}. \tag{5.26}$$

for $N \geq N_6$.

(5.26) combined with (5.22) we get that

$$\begin{aligned}
\mathbb{P}_N\left(BA^2 \neq \emptyset, E\right) &\leq \sum_{v \in B\left(\left\lceil \frac{2\alpha+K+2}{\alpha_3} \right\rceil\right)} \mathbb{P}_N(Z(v) \cap E) \\
&\leq c_2 k^{-1}
\end{aligned} \tag{5.27}$$

with $c_2 = c_2(\lambda_0, \lambda, \alpha_3, \alpha_2, K)$ for $N \geq N_6$. We set θ such that $k = \lfloor 1/2\theta \rfloor > 4c_2/\varepsilon$. A combination of (5.27) and (5.21) gives that

$$\begin{aligned} \mathbb{P}_N\left(BA^2 \neq \emptyset\right) &\leq \mathbb{P}_N\left(BA^2 \neq \emptyset,\, E\right) + \mathbb{P}_N\left(E^c\right) \\ &\leq c_2 k^{-1} + \varepsilon/4 < \varepsilon/2 \end{aligned} \tag{5.28}$$

for $N \geq N' = \bigvee_{i=0}^{6} N_i$.

A proof analogous to that of (5.28) gives that there is $N'' = N''(\alpha, \lambda, K)$

$$\mathbb{P}_N\left(BA^1 \neq \emptyset\right) < \varepsilon/2 \tag{5.29}$$

for $N \geq N''$. A combination of (5.13), (5.28) and (5.29) finishes the proof of Proposition 3.5.

□

A Appendix

A.1 Winding number of arms

Here we prove Proposition 2.7. The proof is motivated by [3]. There, among many other things, it was shown that when there are k disjoint open arms in $A(M, aM)$ $(a > 1)$, then, with conditional probability at least $1 - a^{-\varepsilon}$, and uniformly in M, are also k disjoint open arms which wind around the origin at least $c \log a$ times where c, ε are positive constants.

We prove a slightly different result, namely that if we have k disjoint arms with any colour sequence $\sigma \in \{o, c\}^k$ in $A(M, aM)$, than with conditional probability at least $1 - a^{-\varepsilon}$, these arms wind around the origin at least $c \log a$ times for some $c, \varepsilon > 0$. Following [16], we recall the notion of well separated arms. We modify Definition 7 of [16] for annuli:

Definition A.1. Consider some annulus $A = A(v; M, \tau M)$ and a parallelogram $B = B(v; \tau M)$ for $M \in \mathbb{N}$, $\tau \in (1, \infty)$ and $v \in V$. Let s_T, s_B, s_L, s_R denote the top, bottom, left and right sides of B. Let $\mathcal{C} = \{c_i\}_{1 \leq i \leq j}$ be a set of j disjoint arms in A such that for each i, all of the vertices of c_i are open or all of them are closed. Let z_i be the endpoint of c_i on $\partial B(v; \tau M)$. Let $\eta \in (0, 1]$, we attach a parallelogram r_i to z_i as follows:

$$r_i = \begin{cases} z_i + [-\eta M, \eta M] \boxtimes \left[0, 2\sqrt{\eta} M\right] & \text{if } z_i \in s_T \\ z_i + [-\eta M, \eta M] \boxtimes \left[0, -2\sqrt{\eta} M\right] & \text{if } z_i \in s_B \\ z_i + \left[-2\sqrt{\eta} M, 0\right] \boxtimes [-\eta M, \eta M] & \text{if } z_i \in s_L \\ z_i + \left[0, 2\sqrt{\eta} M\right] \boxtimes [-\eta M, \eta M] & \text{if } z_i \in s_R. \end{cases}$$

We say that $\mathcal{C}$ is η-well-separated on the outside, if the two following conditions are satisfied:

1. The extremities z_i $i = 1, 2, \ldots, j$ are neither too close to each other:

   $$\forall i \neq l,\ d(z_i, z_l) \geq 10\sqrt{\eta} M,$$

   nor too close to the corners Z_l $l = 1, 2, 3, 4$ of B :

   $$\forall i, j,\ d(z_i, Z_l) \geq 10\sqrt{\eta} M.$$

2. Each r_i is crossed vertically when $z_i \in s_T \cup s_B$, and horizontally when $z_i \in s_L \cup s_R$ by some crossing $\tilde{c}_i$ of the same colour as c_i, and

   $$c_i \text{ is connected to } \tilde{c}_i \text{ in } z_i + A\left(1, \sqrt{\eta} M\right).$$

We say that a set $\mathcal{C} = \{c_i\}_{1 \leq i \leq j}$ of disjoint arms in A *can be made η-well-separated* on the outside, if there exists an set $\mathcal{C}' = \{c'_i\}_{1 \leq i \leq j}$ of disjoint arms in A which is η-well-separated on the outside, and c'_i has the same colour and endpoint on $\partial B(v; M)$ as c_i for $i = 1, 2, \ldots, j$.

Similarly to Definition A.1, we define the η-well-separation on the inside. The following statement follows from Lemma 15 of [16].

Lemma A.2. *For $\tau \in (1,\infty)$, and $\delta > 0$, there exists $\eta(\delta) > 0$ such that for any positive integer N, we have*

$$\mathbb{P}_{1/2}\left(\text{any set of disjoint arms in } A(N,\tau N) \text{can be made } \eta\text{-well-separated on the outside}\right) \geq 1-\delta.$$

Moreover, the same statement holds for well separated arms on the inside.

We prove the following proposition.

Proposition A.3. *Let $k, N \in \mathbb{N}$, $a \in (1,\infty)$, and σ a colour sequence of length k. We divide the annulus $A(N,aN)$ into the annuli $A_i = A\left(2^i N, 2^{i+1}N\right)$ for $i = 0,1,\dots,\lfloor \log_2(a)\rfloor - 1$. Let W denote the set of indices i for which all the arms in A_{3i+1} wind around the origin at least once in the counter-clockwise direction for $i = 0,1,\dots,\lfloor \log_2(a)/3\rfloor - 1$. There are positive constants $c = c(k)$, $\varepsilon = \varepsilon(k)$ and $N_0 = N_0(k)$ such that*

$$\mathbb{P}_{1/2}\left(\mathcal{A}_{k,\sigma}(N,aN), |W| \geq c\log_2 a\right) \geq \left(1-a^{-\varepsilon}\right)\pi_{k,\sigma}(N,aN)$$

for all $a \in (1,\infty)$ and $N \geq N_0$.

Remark A.4. Proposition 2.7 follows from Proposition A.3, since $W = \emptyset$ on the event $\mathcal{A}_{k,l,\sigma}(N,aN)$ when $l \geq 1$.

Proof of Proposition A.3. For $a \leq 2$, the statement is trivial. Hence in the rest of the proof we suppose that $a > 2$. Classical RSW techniques [13] give that for all $k \in \mathbb{N}$ there is $\varepsilon_1 = \varepsilon_1(k) > 0$ such that

$$\pi_{k,\sigma}(N,aN) \geq a^{-\varepsilon_1} \tag{A.1}$$

uniformly in $a \geq 2$, $N \geq 1$ and $\sigma \in \{o,c\}^k$.

Let $\eta \in (0,1/10)$. Let IS_i (OS_i) denote the event that any set of disjoint arms of A_i can be made η-well-separated on the inside (outside). Let WS denote the set of indices $i \in \left\{0,1,\dots,\left\lfloor \frac{\log_2 a}{3}\right\rfloor - 1\right\}$ for which OS_{3i} and IS_{3i+2} both hold. Notice that the events $\{i \in WS\}$ for $i = 1,2,\dots,\left\lfloor \frac{\log_2 a}{3}\right\rfloor - 1$ are independent. Moreover, by Lemma A.2, for any $\delta > 0$ there is $\eta(\delta) \in (0,1/10)$ such that

$$\mathbb{P}_{1/2}(i \in WS) \geq 1-2\delta.$$

Combining this with Hoeffding's inequality we get that c_0, δ, η such that

$$\mathbb{P}_{1/2}(|WS| \leq c_0 \log a) \leq a^{-2\varepsilon_1}. \tag{A.2}$$

This and (A.1) gives that

$$\begin{aligned}\mathbb{P}_{1/2}\left(\mathcal{A}_{k,\sigma}(N,aN) \cap \{|WS| > c_0\log(a)\}\right) &\geq \pi_{k,\sigma}(N,aN) - \mathbb{P}_{1/2}(|WS| \leq c_0\log a)\\ &\geq \pi_{k,\sigma}(N,aN) - a^{-2\varepsilon_1}\\ &\geq \left(1-a^{-\varepsilon_1}\right)\pi_{k,\sigma}(N,aN) \end{aligned} \tag{A.3}$$

for all N.

Let us fix an integer $i \in \left\{0,1,\dots,\left\lfloor \frac{\log_2 a}{3}\right\rfloor - 1\right\}$. Condition on the event $\mathcal{A}_{k,\sigma}\left(N,2^{3i+1}N\right) \cap \mathcal{A}_{k,\sigma}\left(2^{3i+2}N,aN\right) \cap \{i \in WS\}$ and on the configuration in $A(N,aN) \setminus A_{3i+1}$. This conditioning gives that all the arms in A_{3i} can be made η-well-separated on the outside, and all the arms in A_{3i+2} can be made η-well-separated on the inside. This imposes some conditions on the configuration in A_{3i+1} : there is a finite collection of disjoint parallelograms in which certain crossing events have to be satisfied. In order to have k arms with colour sequence σ in $A(N,aN)$, it is enough to connect, with the right colour, the k-tuple of parallelograms corresponding to the well separated versions of these arms on the inner parallelogram to those on the outer parallelogram of A_{3i+1}. There might be more than one choice for this pair of k-tuples of parallelograms. In this case we choose a pair in some deterministic way.

We connect the corresponding pairs of parallelograms by disjoint tubes of width $\sqrt{\eta}2^{3i+1}N$ in A_{3i+1} as in the proof of Lemma 4 of [14] (see Figure 9 of [14]), with the difference that these connections are special: We chose these tubes such that each of them winds around the origin at least twice in the counter-clockwise direction. We add an additional tube which avoids the ones above, connects the boundaries of the inner and the outer parallelograms of A_{3i+1} and winds around the origin at least twice in the counter-clockwise direction.

With standard RSW techniques one can show that the probability of the event that the original tubes are crossed in the hard direction by a path with the appropriate colour, and the additional tube is crossed in the hard direction with an open and a closed path is at least $h > 0$. Here $h = h(k, \eta)$ is independent of i, N and the location of the parallelograms we connected. The open and closed crossings of the additional tube forces all the arms of $A(N, aN)$ to wind around the origin in A_{3i+1} at least once in the counter-clockwise direction. Hence the event $\{i \in W\}$ occurs.

Thus the probability of $\{i \in W\}$ conditioned on the event $\mathcal{A}_{k,\sigma} \cap \{i \in WS\}$ and on the configuration in $A(N, aN) \setminus A_{3i+1}$ is at least h. Note that the event $\{i \in W\}$ only depends on the configuration in A_{3i+1}. Hence, when we condition on the event $\mathcal{A}_{k,\sigma}(N, aN)$ and on the realization of WS, the set W stochastically dominates a set Z, where the elements of Z are sampled from WS independently from each other with probability h.

Hence for $c > 0$ we have

$$\begin{aligned}
&\mathbb{P}_{1/2}\left(|W| \geq c\log_2 a \,|\mathcal{A}_{k,\sigma}(N, aN)\right) \geq \mathbb{P}_{1/2}\left(|W| \geq c\log_2 a, |WS| \geq c_0 \log_2 a \,|\mathcal{A}_{k,\sigma}(N, aN)\right) \\
&\quad = \sum_S \mathbb{P}_{1/2}\left(|W| \geq c\log_2 a \,|\mathcal{A}_{k,\sigma}(N, aN), WS = S\right) \mathbb{P}_{1/2}\left(WS = S \,|\mathcal{A}_{k,\sigma}(N, aN)\right) \\
&\quad \geq \sum_S \mathbb{P}_{1/2}\left(|Z| \geq c\log_2 a \,|\mathcal{A}_{k,\sigma}(N, aN), WS = S\right) \mathbb{P}_{1/2}\left(WS = S \,|\mathcal{A}_{k,\sigma}(N, aN)\right), \qquad (A.4)
\end{aligned}$$

where the summation over $S \subseteq \left\{0, 1, \ldots \left\lfloor \frac{\log_2 a}{3} \right\rfloor - 1\right\}$ with $|S| \geq c_0 \log_2 a$. We split this sum in (A.4) depending on the number of elements of S, and we get

$$\begin{aligned}
&\mathbb{P}_{1/2}\left(|W| \geq c\log_2 a \,|\mathcal{A}_{k,\sigma}(N, aN)\right) \\
&\quad \geq \mathbb{P}\left(Y \geq c\log_2 a\right) \sum_{l \geq c_0 \log_2 a} \mathbb{P}_{1/2}\left(|WS| = l \,|\mathcal{A}_{k,\sigma}(N, aN)\right) \\
&\quad = \mathbb{P}\left(Y \geq c\log_2 a\right) \mathbb{P}_{1/2}\left(|WS| \geq c_0 \log_2 a \,|\mathcal{A}_{k,\sigma}(N, aN)\right), \qquad (A.5)
\end{aligned}$$

where Y is a random variable with distribution $Binom\left(c_0 \log_2 a, h\right)$. Using Hoeffding's inequality, we set $c = c(h), \varepsilon_2(h) > 0$ such that

$$\mathbb{P}\left(Y \geq c\log_2 a\right) \geq 1 - a^{-\varepsilon_2}. \qquad (A.6)$$

By substituting (A.6) and (A.3) to (A.5) we get that

$$\mathbb{P}_{1/2}\left(|W| \geq c\log_2 a \,|\mathcal{A}_{k,\sigma}(N, aN)\right) \geq \left(1 - a^{-\varepsilon_1}\right)\left(1 - a^{-\varepsilon_2}\right)$$

for all $a > 2$ and N, which finishes the proof of Proposition A.3. ☐

With suitable adjustments of arguments above, one can show that the following generalization of Proposition 2.7 holds.

Proposition A.5. *For any $k \in \mathbb{N}$, there are positive constants $c = c(k)$, $\varepsilon = \varepsilon(k)$ such that for all $l, l' \in \mathbb{N}$ with $0 \leq l \leq l' \leq k$*

$$\pi_{k,l,\sigma}\left(n_0(k), N\right) \leq cN^{-\varepsilon} \pi_{k,l',\sigma}\left(n_0(k), N\right)$$

uniformly in N and in the colour sequence σ.

A.2 Existence of long thick paths in nice regions

Recall the Definition 4.3 and 4.4. First we prove Lemma 4.5 which is the special case of Lemma 4.6 where C is (a, b)-nice. Then we show how to modify the proof of Lemma 4.5 to get a proof of Lemma 4.6.

Lemma 4.5. *Let $a, b \in \mathbb{N}$ with $a \geq 2000$. Let C be an (a, b)-nice subgraph of $\mathbb{T}$. Then there is a $\lfloor a/200 - 10 \rfloor$-gridpath contained in C with diameter at least $diam(C) - 2b - 2a - 12$.*

Remark A.6. We believe that the constants in Lemma 4.5 are not optimal.

Proof of Lemma 4.5. Recall the lines below Definition 4.4. To prove Lemma 4.5, it is enough to find a path ζ in C such that $diam(\zeta) \geq d - 2b - 2a - 12$ and $\zeta + B(a/100 - 5) \subset C$. We construct ζ by the following strategy.

We put hexagons on the vertices of $\mathbb{T}$ in the following way: The hexagon corresponding to the vertex v is the regular hexagon with side length $1/\sqrt{3}$ centred around v with one of its sides is vertical. These hexagons give a tiling of the plane $\mathbb{R}^2$. Using this tiling, we look at C as the region in $\mathbb{R}^2$ which is the union of the hexagons which are centred around the vertices of C.

Let $x, y \in C$ such that $d(x,y) = diam(C)$. Let $\gamma \subset \mathbb{R}^2$ be a shortest curve connecting x and y in the region C, that is, γ is a continuous map of $[0,1]$ such that 0 is mapped to x and 1 is mapped to y. We get the path ζ from γ as follows. First we cut down two pieces of γ one from its beginning and one from its end. We call the resulting path γ^2. The remaining path is a shortest path. Hence γ^2 turns left (right) at a vertex $z \in \gamma^2$ if close to z on the left (right) side of γ^2 there is a point of ∂C. To avoid ∂C we make a small detour on the right (left) of γ^2 close to z. The path we get from γ^2 with these detours is ζ. We show that ζ indeed satisfies the conditions above, and finish the proof of Lemma 4.5.

We gave a strategy which involved continuous curves and regions in the plane $\mathbb{R}^2$. We adapt it to the triangular lattice in the following precise proof.

Let $x = (x_1, x_2), y = (y_1, y_2) \in C$ such that $d(x,y) = diam(C)$. We further assume that $x_1 < y_1$ and $d(x,y) = y_1 - x_1$. The other case where $d(x,y) = y_2 - x_2$ can be treated similarly. Let $\tilde{\gamma}$ denote a shortest (having the least number of vertices) path connecting x and y in C.

Note that there are $\binom{2n}{n}$ shortest paths between the vertices 0 and $n\underline{e}_1 + n\underline{e}_2$ in $\mathbb{T}$. However, most of them do not follow closely the straight line between the points. Hence $\tilde{\gamma}$ usually does not resemble a shortest continuous curve connecting x and y.

Step 1. *We choose a specific shortest path between x and y.*

For $u, v \in \mathbb{T}$, let $s(u,v)$ denote the line segment connecting u and v in $\mathbb{R}^2$. This segment naturally induces an oriented path $\sigma(u,v)$ in $\mathbb{T}$ as a sequence of the midpoints of the hexagons which are intersected by $s(u,v)$ as we walk along it from u to v. Note that it can happen that the segment $s(u,v)$ contains a side of a hexagon. In this case, we put only one of the neighbouring hexagons to $\sigma(u,v)$. We say that $\sigma(u,v)$ is a triangular grid approximation of the segment $s(u,v)$. $\sigma(u,v)$ is a shortest path between u and v in $\mathbb{T}$.

Recall the notation in Section 4.1. Let $v, u, u' \in \tilde{\gamma}$ with $v \prec u, u'$ and $u \sim u'$. Then for all $w \in \sigma(v,u)$ there is $w' \in \sigma(v,u')$ with $w \sim w'$. Hence for $v \in \tilde{\gamma}$ there are two cases:

- either $\forall u \in \tilde{\gamma}_{v,y} \setminus \{v\}$ we have $\sigma(v,u) \setminus \{v\} \nsim \partial C$, or
- $\exists w = w(v) \in \tilde{\gamma}_{v,y} \setminus \{v\}$ such that $\forall u \in \tilde{\gamma}_{v,w} \setminus \{v, w\}$ we have $\sigma(v,u) \setminus \{v\} \nsim \partial C$, but $\sigma(v,w) \setminus \{v\} \sim \partial C$.

We perform the following procedure. We start at x. If the first case above holds for $v = x$, then we replace $\tilde{\gamma}$ by $\sigma(x,y)$ and finish the procedure. In the second case we replace $\tilde{\gamma}_{x,w(x)}$ by $\sigma(x, w(x))$, and repeat the procedure for $\tilde{\gamma}_{w(x),y}$ starting from $w(x)$. At each step of the procedure, we move at least one vertex further on $\tilde{\gamma}$, hence the procedure terminates in at most $|\tilde{\gamma}|$ steps. Let γ denote the path we get at the end. At each step of the procedure, we make modifications such that the new path is in C and its length is the same as the old path's. Hence $\gamma \subset C$ and $|\gamma| = |\tilde{\gamma}|$.

The construction of γ shows that γ resembles a shortest curve in $\mathbb{R}^2$: It is a sequence of triangular grid approximations of line segments in $\mathbb{R}^2$. We finish Step 1 with the following claim. It is a simple consequence of the definition of γ :

Claim A.7. As we walk along γ, we turn to the left (right) at $v \in \gamma$ if it has a neighbour in ∂C on the left (right) of γ. That is, if $u, v, w \in \gamma$ with $u \prec v \prec w$ and $\sigma(u,v), \sigma(v,w) \subset \gamma$, with $\sigma(u,v) \cup \sigma(v,w) \neq \sigma(u,v)$, then $v \sim \partial C \cap T(u,v,w)$, where $T(u,v,w)$ denotes the triangle spanned by the vertices u, v, w.

Step 2. *We introduce some notation and assign labels to some of the vertices of γ.*

Let

$$ST := \{v = (v_1, v_2) \in V \mid x_1 < v_1 < y_1\}.$$

By possible shortening γ and redefining x and y, we can assume that $\gamma \subset cl(ST)$, $\gamma \cap \partial ST = \{x, y\}$ and $d(x,y) = diam(C)$.

We set $\alpha := \lfloor a/6 \rfloor - 2 > 0$, and define

$$ST^i := \{v = (v_1, v_2) \in V \mid x_1 + b + i\alpha < v_1 < y_1 - b - i\alpha\}$$

for $i \in \{1, 2\}$. Let x^i (y^i) denote the last (first) vertex of γ which is in the half plane $\{v = (v_1, v_2) \in V \mid v_1 \le x_1 + b + i\alpha\}$ ($\{v \mid v_1 \ge x_1 - b - i\alpha\}$). Let $\gamma^i = \gamma_{x^i, y^i}$. Note that $ST^1 \subset ST^2$ and γ^1 is a subpath of γ^2.

Let $i \in \{1, 2\}$. Since γ^i is a shortest path, it is non self-touching. This combined with $\gamma^i \cap \partial ST^i = \{x^i, y^i\}$ we get that γ^i, cuts $cl(ST^i)$ into two connected components. Let ${ST^i}_L$ (${ST^i}_R$) denote connected component $cl(ST^i) \setminus \gamma^i$ which is on the left (right) had side of γ^i as we walk along it.



Figure 8: The path $\gamma_{x,v} \vee \sigma_{v,w} \vee \gamma_{w,y}$, is shorter than γ by at least $\frac{2}{3}\alpha$ vertices.

For $v \in \gamma^2$, we put a label $l(v) \in \{L, R, N, G\}$ as follows. We denote the set of vertices with label $X \in \{L, R, N, G\}$ by γ^2_X. First we define the labels R and L : For $v \in \gamma^2$, we set $l(v) = L$ $(l(v) = R)$ if $ST_L \cap B(v;\alpha) \cap \partial C \neq \emptyset$ $(ST_R \cap B(v;\alpha) \cap \partial C \neq \emptyset)$. To show that the labels L, R are well-defined, we have to check that for $v \in \gamma^2$ at most one of the sets $ST^1{}_L \cap B(v;\alpha) \cap \partial C$ and $ST^1{}_R \cap B(v;\alpha) \cap \partial C$ is non-empty. Since $2\alpha < a$, this follows from Condition 3 of Definition 4.4. Let $\beta := \lfloor \alpha/3 \rfloor$. For $v \in \gamma^2 \setminus \left(\gamma^2_L \cup \gamma^2_R\right)$ we set $l(v) = G$ if $B(v;\beta) \cap \left(\gamma^2_L \cup^2_R\right) = \emptyset$, and $l(v) = N$ otherwise.

We finish Step 2 with the following claim. Since $4\alpha + 2\beta < a$, it is a simple exercise to prove the following claim using Condition 3 of Definition 4.4:

Claim A.8. Let $u \in \gamma^1_L$ and $v \in \gamma^1_R$. Then there is $w \in \gamma^1_G$ which is in between u and v.

Step 3. *For $v \in \gamma^2$, we define two neighbourhoods $\mathcal{F}_v$ and $\mathcal{G}_v$.*

If $l(v) \in \{G, N\}$ then we set $\mathcal{F}_v := B(v;\alpha)$ and $\mathcal{G}_v := B(v;\beta)$.

If $l(v) \in \{L, R\}$, let f^1 (f^2) as the last vertex when we go backwards (forward) from v along γ which is in $B(v;\alpha)$. If it has label L (R) then we define $\mathcal{E}_v$ as the connected component of $B(v;\alpha) \setminus \gamma_{f^1,f^2}$ on the right (left) hand side of γ_{f^1,f^2}. Similarly we define g^1 and g^2 in for the box $B(v;\beta)$, and $\mathcal{G}_v$.

The combination of $4\alpha < a$, Claim A.7 and Condition 3 of Definition 4.4 gives that

$$\left(\gamma_{f^1,g^1} \cup \gamma_{g^2,f^2}\right) \cap B(v;\beta - 1) = \emptyset.$$

Hence we get

Claim A.9. $\mathcal{F}_v \cap B(v;\beta) = \mathcal{G}_v$ for $v \in \gamma^2$.

Step 4. *We investigate the neighbourhood $\mathcal{G}_v$.*

Claim A.10. $\mathcal{G}_v \cap \partial C = \emptyset$ for $v \in \gamma^2$, and $\mathcal{G}_v \cap \gamma^1 = \emptyset$ for $v \in \gamma^2_L \cup \gamma^2_R$.

Proof of Claim A.10. First we show that $\mathcal{G}_v \cap \partial C = \emptyset$ with a proof by contradiction. Suppose that $\mathcal{G}_v \cap \partial C \neq \emptyset$. The definition of labels give that if $\mathcal{G}_v \cap \partial C \neq \emptyset$, then $l(v) = L$ or R. We further suppose that $l(v) = L$. The case where $l(v) = R$ can be treated similarly. We choose w so that it is one of the closest vertices to v among the vertices of $\mathcal{G}_v \cap \partial C$. See Figure 8.

By the definition of the label L, we have that $w \in ST^2_L \cap B(v;\beta)$. Since $w \in \mathcal{G}_v$, i.e. w is on the right hand side of γ_{f^1,f^2} in $B(v;\alpha)$. Hence some subpath of $\gamma^1 \setminus \gamma_{f^1,f^2}$, denoted by ν, has to separate w from v in $\mathcal{F}_v$. Let us walk from v to w on $\sigma(v,w)$, till we hit ν. Let us denote the explored path by $\sigma(v,v')$, where v' is the last point of the exploration. Let γ' be the path we get when we replace the part of γ between v and v' by $\sigma(v,v')$. Consider the case $v' \prec_{\gamma} v$. The other case where $v' \succ_{\gamma} v$ can be treated similarly. The number of vertices of $\sigma(v, v')$ is at most 2β. However, the number of vertices in ν before v' is at least $\alpha - \beta$. Moreover, $\left|\gamma_{f^1, v}\right| \geq \alpha - \beta$. Hence

$$\begin{aligned} |\gamma| - |\gamma'| &\geq 2(\alpha - \beta) - 2\beta \\ &\geq \frac{2}{3}\alpha > 0. \end{aligned} \tag{A.7}$$

The definition of w gives that $\sigma(v, v') \subset C$, thus $\gamma' \subset C$. Hence γ' connects x and y in C and by A.7, it is shorter than γ. This is a contradicts the definition of γ, hence $\mathcal{G}_v \cap \partial C = \emptyset$ for $v \in \gamma^2$.

The proof of $\mathcal{G}_v \cap \gamma^1 = \emptyset$ for $v \in \gamma^2$ is quite similar to the one above, hence we omit it, and finish the proof of Claim A.10 as well as Step 4. ☐

Step 5. *We define the path ζ.*
We set $\varepsilon = \lfloor \beta/4 \rfloor - 2$. For $j \in \{L, R\}$, let

$$U_j := \bigcup_{v \in \gamma_j^2} B(v; \varepsilon). \tag{A.8}$$

$ST_R^2 \setminus U_L \setminus \gamma^2$ ($ST_L^2 \setminus U_R \setminus \gamma^2$) has one infinite connected component which we denote by Z_R (Z_L). Let ζ_j denote the shortest path in $\partial Z_j \cap ST^2$ which connects the left and the right side of ST_2. We orient ζ_L (ζ_R) so that Z_L (Z_R) is on the left (right) hand side. Note that ζ_L, ζ_R are left-right crossings of ST^2.

Note that ζ_L, ζ_R and γ^2 are non self-touching paths. Since Z_R, Z_L and γ^2 are disjoint, γ^2 is sandwiched between $\zeta_L$and ζ_R. Hence $\zeta_L, \zeta_R, \gamma^2$ can have common vertices, but they cannot cross each other. Thus we get the following claim.

Claim A.11. Let $v \in \zeta_L \cap \zeta_R$. Then $v \in \gamma^2$.

Condition 3 of Definition 4.4 implies the following claim.

Claim A.12. Let $v \in \zeta_L \cap \zeta_R$. If w, the next vertex after v on γ^2 exists, then $w \in \zeta_L \cup \zeta_R$.

Let $\vec{G} = \left(\vec{V}, \vec{E}\right)$ be the directed graph induced by the directed paths ζ_L, ζ_R and γ^2.That is $\vec{G} = \left(\vec{V}, \vec{E}\right)$ where $\vec{V} = \zeta_L \cup \zeta_r \cup \gamma^2$, and $(u, v) \in \vec{E}$ if and only if $u, v \in \nu$, $u \sim v$ and $u \prec_{\nu} v$ for some $\nu \in \left\{\zeta_L, \zeta_R, \gamma^2\right\}$. Using the definition of ζ_L and ζ_R it is a simple exercise to show the following claim.

Claim A.13. $\vec{G}$ has no directed loops.

For $j \in \{L, R\}$ and $z \in \zeta_j$ let $n_j(z)$ be the first vertex of $\zeta_j \cap \gamma^2$ after z on ζ_j. That is, $n_j(z) \in \zeta_j \cap \gamma^2$ with $n_j(z) \succeq_{\zeta_j} z$ and if $z' \in \zeta_j \cap \gamma^2$ with $z' \succ_{\zeta_j} z$ then $z' \succeq_{\zeta_j} n_j(z)$. If there is no such vertex, then we set $n_j(z) = \emptyset$.

We define a directed path ζ by the following procedure. Let z_j denote the starting point of ζ_j for $j \in \{L, R\}$. ζ starts at the vertex z defined as

$$z := \begin{cases} z_L & \text{when } n_L(z_L) = \emptyset, \text{ or, when } n_L(z_L) \neq \emptyset \neq n_R(z_R) \text{ and } n_L(z_L) \succeq_{\gamma^2} n_R(z_R) \\ z_R & \text{otherwise.} \end{cases}$$

Suppose that we are at vertex v in ζ. If v is the endpoint of ζ_L or ζ_R, we terminate the procedure. Otherwise, we define the next vertex of ζ, denoted by w, as follows. For $j \in \{L, R\}$, if $v \in \zeta_j$, then v_j denotes the next vertex after v in ζ_j.

- If $v \in \zeta_L \setminus \zeta_R$, then $w = v_L$
- if $v \in \zeta_R \setminus \zeta_L$, then $w = v_R$
- if $v \in \zeta_L \cap \zeta_R$, and if
  - $v_L, v_R \in \gamma^2$, then the definition of ζ_L and ζ_R gives that $v_L = v_R$ and we take $w = v_L = v_R$
  - $v_L \in \gamma^2$, $v_R \notin \gamma^2$, then $w = v_R$
  - $v_R \in \gamma^2$, $v_L \notin \gamma^2$, then $w = v_L$
  - the case $v_L, v_R \notin \gamma^2$ is impossible by Claim A.12.

We finish Step 5 by showing that ζ is well-defined. The definition of ζ shows that if we view ζ as a directed graph, it is a subgraph of $\overrightarrow{G}$. Hence by Claim A.13 ζ has no directed loops. Thus ζ is self avoiding, and the procedure above terminates after finitely many steps, when ζ reaches the endpoint of ζ_L or ζ_R.

Step 6. *We prove the following claim and finish the proof of Lemma 4.5.*

Claim A.14. $\zeta + B(\varepsilon) \subset C$ and $diam(\zeta) \geq d(x,y) - 2b - 4\alpha$.

Proof of Claim A.14. The definition of ζ shows that ζ is a horizontal crossing of ST^2. Hence $diam(\zeta) \geq d(x,y) - 2b - 4a$. We show that for all $v \in \zeta$ we have $v + B(\varepsilon) \subset C$. There are two cases depending on whether v is contained in γ^2.

Case 1: $v \in \zeta \setminus \gamma^2$. Then $v \in \zeta_L \setminus \gamma^2$ or $v \in \zeta_R \setminus \gamma^2$. We assume that $v \in \zeta_L \setminus \gamma^2$. The case where $v \in \zeta_R \setminus \gamma^2$ can be treated similarly. The definition of ζ_L gives that there is $w \in \gamma_R$ such that $v \in (B(w; \varepsilon+1) \setminus B(w;\varepsilon))$ and $B(v;\varepsilon) \cap \gamma_R = \emptyset$. This combined with $4\alpha + 4\varepsilon + 2 < a$ and Condition 3 of Definition 4.4 gives that $B(v;\varepsilon) \cap \left(\gamma_L^2 \cup \gamma_R^2\right) = \emptyset$.

If $\gamma^2 \cap B(v;\varepsilon) \neq \emptyset$, then $\exists u \in \left(\gamma_G^2 \cup \gamma_N^2\right) \cap B(v;\varepsilon)$. Claim A.10 implies that $C \supset \mathcal{G}_u = B(u;\beta) \supset B(v;\varepsilon)$ since $4\varepsilon < \beta$.

If $\gamma^2 \cap B(v;\varepsilon) = \emptyset$, then the definition of w and Claim A.10 shows that $C \supset \mathcal{G}_w \supset B(v;\varepsilon)$ since $2\beta + 2\varepsilon < \alpha$.

Hence $B(v;\varepsilon) \subset C$ in Case 1.

Case 2: $v \in \zeta \cap \gamma^2$. Since $\zeta \subset \zeta_L \cup \zeta_R$, we assume that $v \in \zeta_L$. The case where $v \in \zeta_R$ can be treated similarly. First we show that $v \notin \gamma_L^2 \cap \zeta_L$.

Suppose the contrary, that is $v \in \gamma_L^2 \cap \zeta_L$. Let w be the starting point of the connected component of v in $\gamma^2 \cap \zeta$. By the definition of ζ, $w \in \zeta_L$. Moreover, for w' the vertex right before w on ζ_L, we have $w' \in \zeta_L \setminus \gamma_2$. Hence there is $u' \in \gamma_R^2$ such that $w' \in B(u';\varepsilon+1)$. Since $v \in \gamma_L$ and $u' \in \gamma_R^2$, by Claim A.8 $\exists u \in \gamma_G^2$ which is between u' and v on γ^2. Note that $w' \in \mathcal{G}_{u'}$. By Claim A.10 we have that $\gamma^2_{u',w} \subset \gamma_N^2$. Hence u is between w and v on γ^2. From the definition of w, we get that $u \in \zeta \cap \zeta_L$.

Note that if we show that $u \in \zeta_R$, then we get a contradiction by the definition of ζ. Hence in order to rule out the case $v \in \gamma_L^2 \cap \zeta_L$ it is enough to show that $u \in \zeta_R$.

Suppose the contrary, that is $u \notin \zeta_R$. Recall the definition of U_L from A.8. We introduce a new set of labels on the vertices of U_L as follows. For $q \in U_L$ there is a vertex $r \in \gamma_L$ such that $q \in B(r;\varepsilon)$. We define

$$l'(q) := \begin{cases} B & \text{if } r \prec_{\gamma^2} u \\ A & \text{otherwise.} \end{cases}$$

Since the choice of r above is not necessarily unique, we have to show that $l'(q)$ is well-defined. It can be easily checked by combining Claim A.10, $4\varepsilon + 4 < \beta$ and $u \in \gamma_G^2$. Moreover a similar argument shows that if $q, q' \in U_L$ with $q \sim q'$, then $l'(q) = l'(q')$.

Since γ^2 is non self-touching, $u \in \gamma^2$ is connected to ∞ in ST_R. Since $u \notin \zeta_R$ it is not connected to ∞ in Z_R. Thus there is a path $\nu \subset U_L$ which separates u from ∞ in ST_R. We can choose ν such that it starts and ends at a vertex neighbouring γ^2. By a possible shortening of ν, can assume that if $u' \in \nu$ with $u' \sim \gamma^2$, than u' is either the starting or the endpoint of ν. Let u_1, u_2 be neighbours of the starting point and the endpoint of ν which are in γ^2. The definition of ν gives that u is in between u_1 and u_2 on γ^2. Using Condition 3 of Definition4.4 and that $u \in \gamma_G^2$ it is easy to check that $l'(u_1) \neq l'(u_2)$.

On the other hand, ν is a connected subset of U_L, hence l' is constant on ν. This is a contradiction, thus $u \in \zeta_R$, which in turn shows that $v \in \gamma_L^2 \cap \zeta_L$.

Hence $v \notin \gamma_L^2 \cap \zeta_L$ but $v \in \zeta \cap \gamma^2 \cap \zeta_L$. The definition of ζ_L gives that $v \notin \gamma_R$. Hence $v \in \gamma_N^2 \cup \gamma_G^2$. By Claim A.10 we get $C \supset \mathcal{G}_v = B(v;\beta) \supset B(v;\varepsilon)$, and we are done in Case 2. Since there are no other cases left, the proof of Claim A.14 is finished. ☐

Since $\zeta + B(\varepsilon) \subset C$ and $diam(\zeta) \geq d(x,y) - 2b - 4\alpha$ hence the $\lfloor \varepsilon/2 \rfloor$-gridpath approximation of ζ is contained in C. It has diameter at least $d(x,y) - 2b - 4\alpha - \varepsilon \geq d(x,y) - 2b - 2a - 12$. Since $\varepsilon = \lfloor \beta/4 \rfloor - 2 \geq a/100 - 5$ this concludes the proof of the Lemma (4.5). ☐

We finish the appendix by proving Lemma 4.6.

Lemma 4.6. *Let $a, b, c \in \mathbb{N}$ with $a \geq 2000$. Let C be subgraph of $\mathbb{T}$ which is (a,b)-nice in $B(c)$. Let C' be a connected component of $C \cap B(c)$. Then there is a $\lfloor a/200 - 10 \rfloor$-gridpath contained in C' with diameter at least $diam(C') - 2b - 2a - 12$.*

Proof of Lemma 4.6. Let $x, y \in C'$ with $d(x, y) = diam(C')$, and choose $\tilde{\gamma}$ as one of the shortest paths connecting x, y in C'. From this point on, we can follow the proof of Lemma 4.5 since we will use Condition 3 of Definition 4.4 for pairs of vertices $u, v \in \partial C$ which also contained in $B(c)$. $\square$

References

[1] Michael Aizenmann and David J. Barsky. Sharpness of the phase transition in percolation models. *Communications in Mathematical Physics*, 108(3):489–526, 1987.

[2] David J. Aldous. The percolation process on a tree where infinite clusters are frozen. *Mathematical Proceedings of the Cambridge Philosophical Society*, 128:465–477, April 2000.

[3] Vincent Beffara and Pierre Nolin. On monochromatic arm exponents for 2D critical percolation. *Annals of Probability*, 39:1286–1304, 2011.

[4] Itai Benjamini and Oded Schramm. Private communication with David Aldous, 1999.

[5] Jean Bertoin. *Random fragmentation and coagulation processes*, volume 102 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2006.

[6] Jean Bertoin. Fires on trees. *Ann. Instit. H. Poincaré Probab. Statist.*, 48(4):909–921, 2012.

[7] Philippe Biane, Richard Durrett, and Rick Durrett. Ten lectures on particle systems. In *Lectures on Probability Theory*, volume 1608 of *Lecture Notes in Mathematics*, pages 97–201. Springer Berlin / Heidelberg, 1995.

[8] Rachel Brouwer. *Percolation, forest-fires and monomer dimers (or the hunt for self-organized criticality)*. PhD thesis, Vrije Universiteit, 2005.

[9] Michael Damron, Artëm Shapozhnikov, and Bálint Vágvölgyi. Relations between invasion percolation and critical percolation in two dimensions. *The Annals of Probability*, 37(6):2297–2331, 2009.

[10] Christophe Garban. *Processus SLE et sensibilité aux perturbations de la percolation critique plane*. PhD thesis, Université Paris-Sud XI, 2008. http://www.umpa.ens-lyon.fr/~cgarban/these.pdf.

[11] Christophe Garban, Gabor Pete, and Oded Schramm. The scaling limits of dynamical and near-critical percolation. In perparation.

[12] Christophe Garban, Gábor Pete, and Oded Schramm. Pivotal, cluster and interface measures for critical planar percolation. arXiv:1008.1378, 2010.

[13] Geoffrey Grimmett. *Percolation*. Springer-Verlag, 2nd edition, 1999.

[14] Harry Kesten. Scaling relations for 2D-percolation. *Communications in Mathematical Physics*, 109:109–156, 1987.

[15] Thomas M. Liggett. *Interacting particle systems*. Springer, 2005.

[16] Pierre Nolin. Near-critical percolation in two dimensions. *Electronic Journal of Probability*, 13:1562–1623, 2008.

[17] Balázs Ráth. Mean field frozen percolation. *Journal of Statistical Physics*, 137:459–499, 2009.

[18] Lucio Russo. On the critical percolation probabilities. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 56:229–237, 1981.

[19] Stanislav Smirnov and Wendelin Werner. Critical exponents for two-dimensional percolation. *Mathematical Research Letters*, 8:729–744, 2001.

[20] W. H. Stockmayer. Theory of Molecular Size Distribution and Gel Formation in Branched-Chain Polymers. *Journal of Chemical Physics*, 11:45–55, February 1943.

[21] Jacob van den Berg, Bernardo N.B. de Lima, and Pierre Nolin. A percolation process on the square lattice where large finite clusters are frozen. *Random Structures and Algorithms*, 40:220–226, March 2012.

[22] Jacob van den Berg and Antal A. Járai. The lowest crossing in two-dimensional critical percolation. *The Annals of Probability*, 31(3):1241–1253, 2003.

[23] Jacob van den Berg, Demeter Kiss, and Pierre Nolin. A percolation process on the binary tree where large finite clusters are frozen. *Electron. Commun. Probab.*, 17(2):1–11, 2012.

[24] George W. Wetherill. Comparison of analytical and physical modeling of planetesimal accumulation. *Icarus*, 88(2):336–354, 1990.